

THE REVIEW OF SYMBOLIC LOGIC
Volume 0, Number 0, xxxx 2015

1 INDICATIVE CONDITIONALS, RESTRICTED
2 QUANTIFICATION, AND NAIVE TRUTH

3 HARTRY FIELD

4 **Abstract.** This paper extends Kripke’s theory of truth to a language with a variably strict con-
5 ditional operator, of the kind that Stalnaker and others have used to represent ordinary indicative
6 conditionals of English. It then shows how to combine this with a different and independently
7 motivated conditional operator, to get a substantial logic of restricted quantification within naive
8 truth theory.

9 **§1. Introduction.** The “naive” notion of truth, according to which for each sentence
10 S of our language, the claim that S is true is equivalent to S itself,¹ appears at first blush to
11 be doomed by the Liar paradox and other related paradoxes. But only at first blush: one of
12 the lessons that can be drawn from Kripke 1975 is that naivety in a theory of truth can be
13 retained if one is willing to give up the hegemony of classical logic. There is little reason
14 to doubt the correctness of classical logic as applied to our most serious discourse, e.g. our
15 most serious physical theories. But the semantic paradoxes arise because truth talk gives
16 rise to some anomalous applications (e.g. “viciously self-referential” ones), and it’s rash
17 to assume that classical logic continues to be appropriate to these applications. Maybe we
18 should generalize logic in a way that allows these anomalies to be treated non-classically,
19 while enforcing classicality in situations where anomalies can’t arise. Kripke’s paper, in
20 particular the parts concerning logics based on Kleene valuation schemes, suggests the
21 possibility of naive truth in this setting: in particular, one can have naive truth in a logic
22 that restricts the general application of excluded middle, but which reduces to classical
23 logic in contexts where the anomalies of truth cannot occur.

24 It isn’t *immediately obvious* that the best response to the paradoxes is to abandon the
25 hegemony of classical logic while retaining the hegemony of naive truth—*prima facie*,
26 the reverse seems at least as attractive. But the costs of restricting naive truth turn out to
27 be extraordinarily high,² and so the program of trying to keep it by restricting the scope of
28 classical logic is one well worth pursuing. Kripke 1975 was the first substantial step.³

29 Kripke’s paper by itself shows the possibility of naive truth only for languages of very
30 limited expressive power. The question arises as to how far his ideas can be generalized,

Received: April 27, 2015.

¹ I ignore ambiguities, indexical elements, etc., so as to be able to concentrate on sentence-*types*.
There are subtleties about how best to extend the idea of naive truth to token utterances, but I will
not be concerned with those issues here.

² See Field 2008, Part II, for a review.

³ In Kripke’s paper, and in the present paper too, we keep the classical *structural* rules for validity:
(a) validity is transitive (in the general form given by the Cut Rule), and (b) valid inference is
a relation between a *set* of premises and a conclusion (as opposed e.g. to a multi-set, where the
number of occurrences of the premise matter, as in logics without structural contraction). The use
of substructural logics is unnecessary.

There is also no need to restrict reasoning by cases, or to embrace dialetheism.

1 and on this there has been some progress in recent years. In particular, there are now
 2 techniques for generalizing it to include certain kinds of conditionals (despite the threat
 3 of Curry-like paradoxes)⁴. But one kind of conditional operator that has not been treated
 4 in the literature on naive truth is “variably strict” conditional operators of the sort that have
 5 been discussed by Stalnaker 1968, Lewis 1974, Pollock 1976, Burgess 1981, and many
 6 others. The rough idea of such a conditional is that it is true at a world w if and only if
 7 at all worlds x where its antecedent is true but that are otherwise only minimally different
 8 from w , its consequent is true. (There are different ways of spelling out this rough idea,
 9 depending mostly on the assumptions made about a relation of relative closeness of worlds;
 10 in this paper I’ll adopt a framework, Burgess semantics, that is as neutral as possible about
 11 this.) Variably strict conditionals are clearly non-monotonic (‘If A then C ’ doesn’t imply
 12 ‘If A and B then C ’); from which it pretty much follows that they are non-transitive.⁵
 13 (They are also non-contraposable.) Their non-monotonicity and resulting non-transitivity
 14 make them significantly different from the sort of conditionals heretofore discussed in
 15 the naive truth literature. The early parts of the present paper provide a method (actually
 16 more than one) of extending Kripke’s theory to cover languages with such a variably strict
 17 conditional—including in Section 6 the important case of languages that also have another
 18 conditional operator for restricted quantification.

19 Proponents of variably strict conditionals have divided over how extensive their appli-
 20 cation is. Some, e.g. Lewis, have taken a variably strict operator to model only “counter-
 21 factual” or “subjunctive” conditionals of ordinary language, and have held that “indicative
 22 conditionals” of ordinary language are represented by the familiar ‘ \supset ’. But it’s well known
 23 that understanding ordinary indicatives in terms of ‘ \supset ’ is *prima facie* counterintuitive—
 24 e.g. on that understanding, “If I run for President, I’ll be elected” comes out true, since I’m
 25 resisting all pressure to run—and nowadays it’s more common to think, with Stalnaker,
 26 that the variably strict conditional account is applicable to ordinary indicative conditionals
 27 as well as “counterfactuals”. The first six sections of this paper are neutral on this issue.

28 But I favor the Stalnaker position, and this is relevant to an important application of
 29 the material in the early sections to the logic of restricted quantification, in Section 7.
 30 Restricted quantification poses a serious challenge to naive truth theory. In such a theory
 31 there are already difficulties with properly handling ordinary restricted quantifications
 32 like “Every true sentence in Jones’ book appeared earlier in Smith’s”, but the difficulties
 33 become far greater when one tries to come up with a plausible account of how these interact
 34 with conditionals in a way that validates plausible laws such as “If all A are B and y is
 35 A then y is B ” and “If everything is B then all A are B ”. I’ve addressed this challenge
 36 before (Field 2014), but in a rather *ad hoc* manner; an ultimate goal of this paper is to
 37 answer the challenge without *ad hocness*, by bringing in a more general logic of indicative
 38 conditionals.

39 **§2. Two-valued and three-valued worlds models for the language of indicative con-**
 40 **ditionals.** Let L be a language whose logical primitives are ‘ \neg ’, ‘ \wedge ’, ‘ \forall ’, ‘ $=$ ’, a unary
 41 necessity operator ‘ \Box ’, and a binary conditional operator ‘ \triangleright ’. An additional conditional for
 42 restricted quantification will be added in Sections 5 and 6. For the moment, let’s suppose
 43 that L doesn’t contain “paradox-prone” terms like ‘True’ that will require special treatment.

⁴ See Restall 2007 for a discussion of such paradoxes and of the difficulties that a naive truth theory must overcome if it is to handle them.

⁵ ‘If A and B then A ’ is clearly valid for them, and with it, transitivity would imply monotonicity.

INDICATIVE CONDITIONALS, RESTRICTED QUANTIFICATION, AND TRUTH 3

1 ‘ \triangleright ’ is supposed to represent the indicative and/or counterfactual conditional of English
 2 and be a “variably strict” conditional in the general ballpark of Lewis, Stalnaker, Pollock
 3 and Burgess. Of these semantics, Burgess’s is the most general (that is, the others can be
 4 obtained by adding restrictions to it),⁶ and I will consider both it and a slight modification
 5 of it. Both versions of the Burgess semantics are initially based on “2-valued worlds
 6 models”, which I’ll now describe. (For simplicity I’ll assume that L has no individual
 7 constants or function symbols; also, that its only variables are first order.)

8 A 2-valued worlds model M for L consists of

9 (i): A non-empty set W of worlds, perhaps with a distinguished non-empty subset
 10 $NORM$ of “normal” worlds. (Nothing central to this paper depends on allowing
 11 non-normal worlds; I do so simply for added generality. The definition of validity
 12 will be in terms of the normal worlds only, but allowing for non-normal worlds
 13 may affect which conditionals can be true at normal worlds.)

14 (ii): For each $w \in W$, a subset W_w of W and a pre-order (reflexive and transitive
 15 relation) \leq_w on W_w .⁷ (Think of W_w as the set of worlds “accessible from” w , and
 16 ‘ $x \leq_w y$ ’ as meaning “the change from w to x is no greater than the change from
 17 w to y ”.)

18 (iii): For each $w \in W$, a non-empty set U_w (the universe of w). Let U be the union of
 19 the U_w .

20 (iv): For each $w \in W$ and k -place predicate p , a function p_w from U^k (the set of
 21 k -tuples of members of U) to $\{0, 1\}$. (The set of k -tuples that get assigned value 1
 22 is the *extension of p* in the model.) We require that the function $=_w$ (associated
 23 with ‘=’) assigns 1 to $\langle o, o \rangle$ for each $o \in U$ and assigns 0 to all other pairs.

24 (W , $NORM$, etc. can all vary from one model to another, so we should really write W_M ,
 25 $NORM_M$, $W_{M,w}$, $\leq_{M,w}$, $U_{M,w}$ and $p_{M,w}$.) Regarding (iv), we could if we like impose
 26 the (“actualist”) requirement that p_w never assign value 1 to k -tuples not in U_w^k ; it won’t
 27 matter for what follows.⁸

28 Regarding (ii), we could if we like impose additional conditions on W_w and \leq_w for each
 29 $w \in W$, or at least for each w in $NORM$. (The distinction of non-normal worlds from
 30 normal ones only matters if some such additional conditions apply only to normal worlds.)
 31 Indeed, one such condition is almost universally regarded as appropriate for indicative and
 32 counterfactual conditionals (at least for worlds w in $NORM$):

33 **Weak Centering:** $w \in W_w$, and for any x in W_w , $w \leq_w x$

34 That Weak Centering holds at least for worlds in $NORM$ is required if Modus Ponens for \triangleright
 35 is to be valid, on the account of validity soon to be given, which involves preservation of

⁶ Not every defensible model of conditionals can be fit into the Burgess framework (or the slight modification of it to be mentioned soon). I suspect that the basic ideas of this paper can be adapted to plausible alternative models, but will not attempt to prove this.

⁷ An alternative convention is to take \leq_w to be a pre-order on the full W , and subject to the constraint that if $y \in W_w$ and $x \leq_w y$ then $x \in W_w$.

⁸ In the 3-valued context to be introduced shortly, we could introduce a more thorough actualism, in which the p_w never assign value 0 or 1 to such k -tuples; in effect this would make U_w^k rather than the full U^k the domain of p_w . But again, this would make no difference to the issues I’m concerned with.

1 value 1 at normal worlds.⁹ (Modus Ponens has been questioned for indicative conditionals
2 (McGee 1985), but the grounds for doing so seem weak in the context of the semantics for
3 variably-strict conditionals.)¹⁰

4 In addition to Weak Centering, Lewis, Stalnaker, Pollock and many others also accept
5 one or more of the following conditions (for all worlds or just for normal ones):

6 **Strong Centering:** $w \in W_w$, and for any x in W_w other than w , $w <_w x$ (i.e. $w \leq_w x$
7 and not($x \leq_w w$))

8 **No Incomparabilities:** for any x, y in W_w , either $x \leq_w y$ or $y \leq_w x$

9 **No Ties:** for any distinct x, y in W_w , not both $x \leq_w y$ and $y \leq_w x$

10 **Limit Condition:** the relation $<_w$ is well-founded.

11 What follows will be completely neutral as to which if any such conditions are imposed,
12 except for occasional reminders that restricting to models with Weak Centering (at least at
13 normal worlds) is advantageous.¹¹

14 To simplify the presentation of the semantics I adopt the usual trick of expanding the
15 language to contain a new name for each object in U ; call the expanded language L^+ . (The
16 expansion depends on the underlying model, so we should really write L_M^+ .) I'll consider
17 two ways of evaluating the sentences of L^+ in M .

18 The first version is 2-valued:

19 **Burgess evaluation procedure:**

- 20 • $|p(c_1, \dots, c_k)|_w$ is just $p_w(o_1, \dots, o_k)$, where c_1, \dots, c_k are the names for o_1, \dots, o_k
21 respectively.

⁹ Demanding Weak Centering at non-normal worlds as well as normal ones would lead in addition, in the current 2-valued framework, to the validity of the inference from $C \triangleright A$ and $C \triangleright (A \triangleright B)$ to $C \triangleright B$. If we want Modus Ponens without getting that even for 2-valued sentences, we need the flexibility provided by non-normal worlds. In general, the point of non-normal worlds is to provide such added flexibility as to what comes out valid.

I've said that nothing in this paper depends on making use of such added flexibility: there will be no need to have the flexibility in the logic that includes 'True' if one doesn't utilize it in the base logic without 'True'. This may seem surprising: we presumably want Modus Ponens for \triangleright , but we don't want the law just cited since by taking A to be C we'll be led to the inference from $C \triangleright (C \triangleright B)$ to $C \triangleright B$, which in combination with Modus Ponens is well known to rule out naive truth by Curry's paradox. But there is actually no problem: in the semantics to be introduced, Weak Centering at *all* worlds guarantees only that the inference from $C \triangleright A$ and $C \triangleright (A \triangleright B)$ to $C \triangleright B$ will hold for 2-valued sentences; and the sentences involved in Curry-type paradoxes will not be 2-valued. (Modus Ponens, on the other hand, will be guaranteed for *all* sentences, even by Weak Centering just at normal worlds.)

¹⁰ The canonical supposed counterexample involves a 3-candidate race whose leading candidates are a Democrat and a Republican, with an Independent far behind. Then the claim "If the Republican doesn't win, the Independent will" seems false. But "The Democrat won't win" may be true, and "If the Democrat doesn't win, then if the Republican doesn't win the Independent will win" may seem true; and these two claims lead to the false claim by Modus Ponens. A standard resolution of this, which I support, is that the complex conditional that "may seem true" isn't: what's true is only that if the Democrat doesn't win *and* the Republican doesn't win then the Independent will win, but to get from that to the complex conditional one needs the rule of Exportation $(A \wedge B) \triangleright C \models A \triangleright (B \triangleright C)$, which is invalid on the variably-strict semantics.

¹¹ It's also possible to add "purely modal" conditions, not involving the \leq_w ; e.g.

S4 if $x \in W_w$ and $y \in W_x$ then $y \in W_w$.

What follows is neutral on these as well.

- 1 • $|\neg A|_w$ is $1 - |A|_w$
2 • $|A \wedge B|_w$ is $\min\{|A|_w, |B|_w\}$
3 • $|\forall x A|_w$ is $\min\{|A(c/x)|_w : \text{all } c \text{ that name members of } U_w\}$
4 • $|\Box A|_w$ is $\min\{|A|_x : x \in W_w\}$
5 • $|A \triangleright B|_w = \begin{cases} 1 & \text{iff } (\forall x \in W_w)[|A|_x = 1 \supset \\ & (\exists y \leq_w x)[|A|_y = 1 \wedge (\forall z \leq_w y)(|A|_z = 1 \supset |B|_z = 1)]] \\ 0 & \text{iff } (\exists x \in W_w)[|A|_x = 1 \wedge \\ & (\forall y \leq_w x)[|A|_y = 1 \supset (\exists z \leq_w y)(|A|_z = 1 \wedge |B|_z = 0)]] \end{cases}$

6 (Let a w -neighborhood be a non-empty subset N of W_w such that if $x \in N$ and $y \leq_w x$
7 then $y \in N$; and call a w -neighborhood A -consistent if it contains a world where $|A|$ is 1.
8 Then the right hand side of the 1-clause for \triangleright says that all A -consistent w -neighborhoods
9 have A -consistent sub- w -neighborhoods throughout which if $|A|$ is 1, so is $|B|$; and the
10 right hand side of the 0-clause says that there is an A -consistent w -neighborhood such
11 that every A -consistent sub- w -neighborhood of it contains a world where $|A|$ is 1 and $|B|$
12 is 0. If one were to make the ‘‘No Incomparabilities’’ assumption (for all worlds, not just
13 normal ones) one could simplify these clauses for \triangleright a bit: that assumption amounts to the
14 assumption that for each w , the w -neighborhoods are nested; and given that, the 1-clause is
15 equivalent to the claim that if there is at least one A -consistent w -neighborhood then there
16 is one throughout which if $|A|$ is 1, so is $|B|$.)

17 These stipulations give every L^+ -sentence a unique value in $\{0,1\}$ at each world, given
18 any 2-valued worlds model M . Conditionals don’t in general contrapose, but they shouldn’t:
19 ‘If Trump runs for President he won’t be elected’ shouldn’t imply ‘If Trump is elected he
20 won’t have run’.

21 Validity is explained as follows:

22 **(VAL):** An inference from a set Γ of L -sentences to an L -sentence B is *Burgess-valid* if
23 for every worlds model M and every $w \in NORM_M$, if $|A|_{M,w} = 1$ for all A in Γ then
24 $|B|_{M,w} = 1$.

25 (Here what counts as a worlds model depends on which structural conditions (e.g. Weak
26 Centering) have been imposed, so (VAL) really gives a family of notions of validity. Again,
27 the restriction to normal worlds only makes a difference when one imposes structural
28 requirements on the normal worlds of models that don’t apply to all worlds.)¹²

29 We define \vee from \wedge and \neg , and \exists from \forall and \neg , and \diamond from \Box and \neg , in the usual ways.
30 ($|\diamond A|_w$ is thus $\max\{|A|_x : x \in W_w\}$.)¹³ $A \triangleleft B$ will abbreviate $(A \triangleright B) \wedge (B \triangleright A)$.

31 But there might be a reason to treat ‘ \triangleright ’ slightly differently. Many people, myself in-
32 cluded, find it natural to suppose that $\neg(A \triangleright B)$ should be equivalent to $A \triangleright \neg B$, modulo

¹² We can generalize to the case where B and the members of Γ can contain free variables: for any
model M , if f is any function assigning $\{L^+\}_M$ -names to free variables, and A is any L -formula,
let A^f be the $\{L^+\}_M$ -sentence that results from substitution by f . Then the generalization of
(VAL) is

(VAL_{gen}): An inference from a set Γ of L -formulas to an L -formula B is *Burgess-valid* if for
every worlds model M and every f for M and every $w \in NORM_M$, if $|A^f|_{M,w} = 1$ for all
 A in Γ then $|B^f|_{M,w} = 1$.

¹³ Why take ‘ \Box ’ as primitive, since $\Box A$ is equivalent to $\neg A \triangleright A$? The answer is that the equivalence
will be lost once we move to a 3-valued semantics, either because of the move to the modified
Burgess evaluation procedure to be given next or to handle predicates like ‘True’.

1 the assumption $\diamond A$ (that is, each should imply the other on that assumption). We don't
 2 have that on the above semantics, unless we add strong assumptions (viz.: No Ties, No
 3 Incomparabilities and the Limit Condition); that was one of Stalnaker's arguments for
 4 imposing those assumptions. If we want that equivalence without the strong assumptions,
 5 we can get it by strengthening the 0 clause for ' \triangleright ' while leaving the 1 clause as is. We then
 6 need a 3-valued framework to handle sentences that receive neither value 1 nor value 0.
 7 Our worlds models are still 2-valued for the moment, i.e. atomic sentences of L^+ can only
 8 take values in $\{0,1\}$, but we allow an additional value $\frac{1}{2}$ for conditionals and sentences
 9 containing them as components. The evaluation clause for ' \triangleright ' is as follows:

10 **Modified Burgess evaluation procedure**¹⁴

$$11 \quad |A \triangleright B|_w = \begin{cases} 1 & \text{iff } (\forall x \in W_w)[|A|_x = 1 \supset (\exists y \leq_w x)[|A|_y = 1 \wedge \\ & (\forall z \leq_w y)(|A|_z = 1 \supset |B|_z = 1)]] \\ 0 & \text{iff } (\forall x \in W_w)[|A|_x = 1 \supset (\exists y \leq_w x)[|A|_y = 1 \wedge \\ & (\forall z \leq_w y)(|A|_z = 1 \supset |B|_z = 0)] \wedge (\exists x \in W_w)(|A|_x = 1) \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

12 (The 0-clause says that there are A -consistent w -neighborhoods, and each such has A -
 13 consistent sub- w -neighborhoods throughout which if $|A|$ is 1 then $|B|$ is 0.) I've already
 14 written the evaluation clauses for \neg , \wedge , \forall and \Box in a way that carries over automatically to
 15 allow for the extra value. (These clauses are called the *Strong Kleene rules*.)

16 The crucial thing about this alternative evaluation procedure for \triangleright is that if $|\diamond A|_w$ is 1,
 17 i.e. if $(\exists x \in W_w)(|A|_x = 1)$, then $|\neg(A \triangleright B)|_w$ is just $|A \triangleright \neg B|_w$. Of course, a consequence
 18 will be a minimal non-classicality: excluded middle can fail for sentences containing ' \triangleright '.
 19 The cost of this isn't that high, I think: indeed, once we introduce a truth predicate, we'll
 20 need excluded middle to fail even more broadly than that.

21 What notion of validity goes with this modified evaluation scheme? There are several
 22 possible choices, but the one I will work with carries over the wording of (VAL) (or more
 23 generally, the (VAL_{gen}) of note 12) to the 3-valued case: validity involves preservation
 24 of value 1 at all normal worlds in all models (with the values now given by the modified
 25 evaluation rules).

26 In adding 'True' to the language we will need to adapt either the original Burgess se-
 27 mantics or the modified Burgess semantics to 3-valued worlds models. A *3-valued worlds*
 28 *model* is just like a 2-valued one except that in clause (iv) we replace $\{0, 1\}$ with $\{0, \frac{1}{2}, 1\}$,
 29 so that atomic sentences as well as conditionals can receive value $\frac{1}{2}$.¹⁵ So 2-valued models
 30 are a special case of 3-valued. The most straightforward adaptation to the presence of
 31 'True' would be to simply use the Burgess or modified Burgess rules as written. But instead
 32 of doing precisely that I will proceed in a more roundabout way, which nonetheless is
 33 modeled on these rules and agrees with them entirely for conditionals whose antecedent
 34 and consequent don't contain 'True'.

¹⁴ There's no danger of this requiring that the same conditional get both value 0 and 1 at a world. For
 assume as an induction hypothesis that A and B each have a unique value at each world. (Actually
 we'll only need that B does.) If $A \triangleright B$ gets value 0 at w , then there must be a $y \in W_w$ for which
 $|A|_y$ is 1 and $(\forall z \leq_w y)(|A|_z = 1 \supset |B|_z = 0)$; and if it gets 1 there must be a $y^* \leq_w y$ such
 that $|A|_{y^*} = 1 \wedge (\forall z \leq_w y^*)(|A|_z = 1 \supset |B|_z = 1)$. But these require that $|B|_{y^*}$ is both 0 and
 1, contrary to the induction hypothesis.

¹⁵ For present purposes I keep the earlier restriction on the assignment $=_w$ to '=', though with the
 third value it could be liberalized somewhat to allow for indeterminate identity.

1 Validity will be defined as before: preservation of value 1 at all normal worlds of all
 2 models (that meet whatever structural conditions such as Weak Centering that one has
 3 imposed). However, when L contains a truth predicate we'll restrict the models used in the
 4 definition, to "arithmetically standard" models that treat the predicate 'True' in a certain
 5 way. The details are in Section 3.

6 **§3. Truth and satisfaction: the strategy.** Suppose that L contains a truth predicate
 7 (more specifically, a predicate of truth in L).¹⁶ To be of interest, L will also need to have
 8 the resources to talk of the bearers of truth, i.e. sentences, and their syntactic properties. Or
 9 instead of syntactic objects, L could just contain arithmetic; we could talk of truth relative
 10 to a Gödel numbering. A language with a satisfaction predicate (from which truth can be
 11 defined, but not in general conversely) is more interesting; but to have a useful satisfaction
 12 predicate we need to be able talk of finite sequences of arbitrary objects from the universe
 13 of discourse, which requires additional mathematical resources. Moreover, dealing with
 14 satisfaction involves some notational complexity that can be confusing. So to keep things
 15 simple I'll take L to involve a truth predicate but not a satisfaction predicate. It is routine to
 16 generalize what follows from truth to satisfaction (when the extra mathematical resources
 17 are available in L).

18 Rather than building syntactic notions into L , I'll follow the Gödel numbering route: L
 19 will contain the predicates 'natural number', 'is zero', 'is the successor of', 'is the product
 20 of', and '='. (I'll fix a Gödel numbering g of L .) I'll also be concerned only with worlds
 21 models M whose arithmetic part is standard (an ω -model) and the same from world to
 22 world. That is, I'll assume that in every model and every world in it, U_w is a superset
 23 of the set N of natural numbers, and 'natural number' is assigned N as its extension,
 24 and the other arithmetic vocabulary is interpreted in the standard way. I'll call worlds
 25 models meeting these restrictions *arithmetically standard*. It's natural to restrict to them
 26 since without some such restriction the Gödel numbering results in "non-standard syntactic
 27 expressions" that have infinitely many distinct sub-expressions. If in defining validity we
 28 restrict to arithmetically standard worlds models, the result is ω -validity (or *validity in*
 29 *ω -logic*); it is this rather than regular validity that I will be primarily concerned with.

30 Kripke 1975, at least the part dealing with the Kleene construction, was concerned with
 31 the possibilities for *naive* truth (and satisfaction), though in languages not containing \triangleright .
 32 Here I will extend his results to languages containing \triangleright .

33 I informally defined "naive theory of truth" in my introductory remarks, but I should
 34 be more precise. Let a formula Y be a *Tr-equivalent* of a formula X if there are (possibly
 35 multiple) L -sentences A such that Y results from X by (possibly multiple) substitutions of
 36 $True((A))$ for A and/or vice versa. A *naive theory of truth* is one where whenever Y is a
 37 Tr-equivalent of X , Y follows from X and vice versa (i.e. the inferences from X to Y and
 38 Y to X are valid). The semantic paradoxes show that naivety is unattainable in classical
 39 logic, but Kripke (in his Kleene-based construction) showed it attainable in non-classical
 40 logic, by the use of 3-valued models. (Again, his language didn't contain \triangleright .)

41 Naivety is not the sole requirement we should impose on a theory of truth: we also want
 42 it to obey reasonable compositional laws, and to allow the truth predicate to appear in an
 43 induction rule. More on these shortly.

¹⁶ Not in L^+ : the new names in L^+ aren't part of the language L for which we're giving a truth theory, and are dependent on a particular model of L . Any apparent loss in restricting truth to L -sentences should be met by generalizing from truth to satisfaction, as discussed later in the paragraph.

1 Our theory of truth should of course also be consistent, at least Post-consistent: that
 2 is, it shouldn't imply everything. I don't *in principle* require negation-consistency, i.e. the
 3 restriction to theories that for no A imply both A and $\neg A$. However, as is implicit in my
 4 earlier definition of validity, the theories I'll be developing satisfy disjunctive syllogism
 5 ($A \vee B, \neg A \models B$), and for those theories Post-consistency requires negation-consistency.
 6 (While there are familiar "paraconsistent" logics that avoid paradoxes without restricting
 7 excluded middle, by restricting disjunctive syllogism instead, they don't seem to me a
 8 promising framework for my ultimate goal of restricted quantification: the comments in
 9 Section 7 below on Beall *et al* 2006 and Beall 2009 may be enough to give some sense
 10 of this.)

11 Actually we want our naive truth theory to be more than (Post- or negation-) consistent:
 12 a consistent theory might, after all, imply the defeat of the Paris Commune, and no logic
 13 of truth should do that. What we want is for our theory of truth to be "consistent with any
 14 arithmetically standard worlds model" of the 'True'-free fragment of L , which I'll call L_0 .
 15 More fully,

16 **GOAL:** We want to generate from each 2-valued arithmetically standard worlds model
 17 M_0 for L_0 a corresponding 3-valued worlds model M for L that (a) validates naive truth
 18 and (b) is exactly like M_0 except that it assigns a 3-valued extension to 'True'. It follows
 19 from (b) that the sentences of L_0^+ get the same value at w in M as in M_0 , for each world
 20 w ; and also that M is arithmetically standard, given that M_0 is.

21 I'll take the allowable worlds models M of L to be just the ones generated from worlds
 22 models M_0 of L_0 in this way; that is, validity, consistency etc. *in the logic of truth* are
 23 defined by quantification over the arithmetically standard worlds models M_0 of the 'True'-
 24 free fragment of the language, and extending the valuation to sentences with 'True' by a
 25 procedure to be given.¹⁷ (It isn't immediately obvious what this procedure should be when
 26 it comes to sentences containing both ' \triangleright ' and 'True': e.g. to take a very simple Curry-like
 27 case, it isn't immediately obvious how to evaluate a sentence K_{\triangleright} constructed by the usual
 28 Gödel-Tarski techniques so as to be equivalent to $True(\langle K_{\triangleright} \rangle) \triangleright \neg True(\langle K_{\triangleright} \rangle)$). Indeed I
 29 will consider several alternative procedures for constructing the extension.)

30 Note that if we can establish (GOAL), we get a kind of conservativeness result: letting
 31 *-consistency be consistency in ω -logic, we have that any classically *-consistent set of
 32 sentences of L_0 is *-consistent in a naive truth theory.¹⁸ The naive truth theory in question
 33 includes not merely the inferences from any sentence to its Tr-equivalents, it can include
 34 any other law validated in the construction of M from M_0 . What laws these are will of
 35 course depend on the details of the construction of M from M_0 , which is yet to be given.

36 But whatever the details, it is clear in advance that if (GOAL) is achieved then our
 37 construction will not only be one on which truth is naive, but one where mathematical
 38 induction in the form $A(0) \wedge (\forall n \in N)(A(n) \supset A(n+1)) \models (\forall n \in N)A(n)$ is legitimate
 39 even for formulas containing 'True'.¹⁹ The reason is that in any arithmetically standard

¹⁷ A slightly more general procedure will be mentioned in note 30.

¹⁸ Calling this a conservativeness result could be misleading: there is no *deductive* conservativeness, it is a kind of semantic conservativeness in ω -logic. Its purpose, as I've said, is to ensure that the set of principles to be declared valid in the naive truth theory is not merely consistent, but consistent with any set of assumptions in the 'True'-free language that are compatible with the conditional logic and standard models of arithmetic.

¹⁹ Analogous forms with other modus-ponens-obeying conditionals in place of the ' \supset ' are guaranteed too.

1 worlds model, when the premises of this induction rule hold at a world the conclusion must
2 too, and the construction guarantees that the new worlds model is arithmetically standard.

3 It is almost as immediate that the construction will validate the desirable composition
4 principles, e.g.

5 **COMPOS-GENERAL:** $\forall x \forall y \forall z$ (If x and y are sentences and z is the result of applying
6 ‘ \triangleright ’ to x and y in that order, then $\Box [True(z)$ if and only if $(True(x) \triangleright True(y))$]).

7 For as long as the logic validates each instance of “ $\Box [A$ if and only if $A]$ ”, then the naivety
8 of truth guarantees the validity of each instance of

9 **COMPOS-SCHEMA:** $\Box [True((A \triangleright B))$ if and only if $(True((A)) \triangleright True((B)))$];

10 and since the constructed model is arithmetically standard, the generalization is guaranteed
11 to hold in the model when the instances do. [This holds on *any* reading of ‘if and only if’,
12 as long as “ $\Box [A$ if and only if $A]$ ” is validated. At the moment, the only available reading
13 is ‘ \triangleleft ’, but I will later add other biconditionals, and the point applies equally to them.]

14 **§4. Truth and satisfaction: the details.** I now outline a generalization of Kripke’s
15 construction. The initial generalization, which takes ‘ \triangleright ’ as a black-box, is completely
16 routine, hardly a generalization at all; but a non-Kripkean ingredient is then required, to
17 give a substantial account of ‘ \triangleright ’.

18 Let’s get the pure Kripke part of the construction out of the way first. It’s clear from
19 what has already been said that each of the worlds w in the model for L will be evaluated
20 in part on the basis of U_w and the w -extensions of L_0 -predicates. The additional ingredients
21 needed to evaluate L^+ -sentences at each w are:

- 22 • a 3-valued extension T_w for ‘True’: it assigns values in $\{0, \frac{1}{2}, 1\}$ to objects in
23 U . (We’ll want it to assign non-zero values only to those objects that are Gödel
24 numbers of L -sentences under the chosen Gödel numbering.)
- 25 • a function j_w that assigns to each L^+ -sentence of form ‘ $A \triangleright B$ ’ a value in $\{0, \frac{1}{2}, 1\}$.

26 Let T and j be the functions that assign to each $w \in W$ a T_w and j_w . Relative to any such
27 T and j , the Kleene rules tell us how to evaluate every L^+ -sentence at w :

- 28 • For p other than ‘True’, $|p(c_1, \dots, c_k)|_{w,j,T}$ is just $p_w(o_1, \dots, o_k)$;
- 29 • $|True(c)|_{w,j,T}$ is $T_w(o)$, where o is the object denoted by the L^+ -name c ;
- 30 • $|\neg A|_{w,j,T}$ is $1 - |A|_{w,j,T}$
- 31 • $|A \wedge B|_{w,j,T}$ is $\min \{|A|_{w,j,T}, |B|_{w,j,T}\}$
- 32 • $|\forall x A|_{w,j,T}$ is $\min \{|A(c/x)|_{w,j,T} : \text{all } c \text{ that name members of } U_w\}$
- 33 • $|\Box A|_{w,j,T}$ is $\min \{|A|_{w,j,T} : x \in W_w\}$
- 34 • $|A \triangleright B|_{w,j,T} = j_w(A \triangleright B)$.

35 The important thing about this is a monotonicity principle. Let $T \leq_K T^*$ mean that for
36 every w and every L -sentence S , if $T_w(S) = 1$ then $T_w^*(S) = 1$ and if $T_w(S) = 0$ then
37 $T_w^*(S) = 0$. Then

38 **(MONOT):** For any M and j : if $T \leq_K T^*$ then for any $w \in W$ and any L^+ -sentence A ,
39 if $|A|_{w,j,T} = 1$ then $|A|_{w,j,T^*} = 1$ and if $|A|_{w,j,T} = 0$ then $|A|_{w,j,T^*} = 0$.

40 This is easily proved by an induction on the complexity of A . (The result is familiar from
41 Kripke 1975, except that I’ve added a trivial \triangleright clause and a world-argument for T .)

42 This is the background for

1 PROPOSITION. [*Kripke's observation.*] For any M and j , there are T ("Kripke fixed
2 points" relative to M and j) for which, for each $w \in W$:

3 For every L -sentence A , $|A|_{w,j,T} = T_w(g(A))$ [and hence $|A|_{w,j,T} = |True(c)|_{w,j,T}$,
4 where c denotes $g(A)$]; and

5 $T_w(o)$ is 0 if o is not $g(A)$ for some L -sentence A .

6 In particular, for any M and j there is a minimal fixed point T_{min} , i.e. a fixed point (relative
7 to M and j) such that for every other fixed point T (relative to M and j), $T_{min} \leq_K T$.

8 Kripke's observation is easily proved by transfinite induction.²⁰

9 It easily follows that as long as j is transparent, in the sense that it assigns Tr-equivalent
10 formulas the same value, then the naivety condition is met: whenever A and B are Tr-
11 equivalent, $|A|_{w,j,T_{min}} = |B|_{w,j,T_{min}}$. (And similarly for fixed points T other than T_{min} .)

12 The definition of T_{min} depended on the choice of M and j , but given those, T_{min} is
13 uniquely determined; so we can abbreviate $|A|_{w,j,T_{min}}$ as $|A|_{w,j}$. To repeat, this valuation
14 yields naive truth as long as j is transparent.

15 The harder task is to construct an appropriate transparent j -function for evaluating
16 conditionals at worlds. What we want is a transparent j that leads to a logic that reduces
17 to the Burgess or modified-Burgess logic when applied to 'True'-free sentences and which
18 weakens the laws as little as possible when sentences with 'True' are allowed as instances.
19 There are at least two approaches to constructing such a j function: a revision construction,
20 with similarities to those in Field 2008; or a fixed point construction, with similarities to
21 those in Field 2014.

22 The revision construction is simpler, so I'll focus on it, but will also make a few remarks
23 about the (perhaps more aesthetically pleasing) fixed point construction.

24 **4.1. The revision construction.** Fix a worlds model M_0 for L_0 . Suppose we have given
25 a provisional valuation j_v , which assigns values $|B \triangleright C|_{w,j_v}$ to any L^+ -sentences B and C .
26 As we've seen, this indirectly gives a value $|A|_{w,j_v}$ to every L^+ -sentence A at every world,
27 via the Kripke minimal fixed point construction; let's just write this as $|A|_{w,v}$. We want
28 to use this valuation j_v to construct a revised one j_{v+1} , perhaps a better one, which is
29 transparent if the original one is; the structure of worlds is used in the revision.

30 There are two possibilities for j_{v+1} , one based on the original Burgess valuation rules
31 and the other based on the variant. For the original it is:

$$32 \quad j_{w,v+1}(A \triangleright B) \text{ is } \begin{cases} 1 & \text{iff } (\forall x \in W_w)[|A|_{x,v} = 1 \supset (\exists y \leq_w x)[|A|_{y,v} = 1 \wedge \\ & (\forall z \leq_w y)(|A|_{z,v} = 1 \supset |B|_{z,v} = 1)]] \\ 0 & \text{iff } (\exists x \in W_w)[|A|_{x,v} = 1 \wedge (\forall y \leq_w x)[|A|_{y,v} = 1 \supset \\ & (\exists z \leq_w y)(|A|_{z,v} = 1 \wedge |B|_{z,v} = 0)]] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

33

²⁰ Holding M and j fixed, we define T_0 to be the function assigning the value $\frac{1}{2}$ to every Gödel
number of an L -sentence, and 0 to everything else; $T_{\sigma+1}$ the function assigning every world w
and L -sentence A the value $|A|_{w,j,T_\sigma}$; and T_λ (for limit λ) the function assigning every world w
and L -sentence A the value

$$\begin{cases} 1 & \text{if for some } \sigma < \lambda \text{ and every } \tau \text{ such that } \sigma \leq \tau < \lambda, |A|_{w,j,T_\tau} = 1; \\ 0 & \text{if for some } \sigma < \lambda \text{ and every } \tau \text{ such that } \sigma \leq \tau < \lambda, |A|_{w,j,T_\tau} = 0; \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

We can then easily prove by induction that if $\sigma < \tau$, $T_\sigma \leq_K T_\tau$. Cardinality considerations
then show that there are ordinals σ (of the cardinality of U_M) after which the assigned T never
changes. Taking T_{min} to be T_σ for such a σ , we get the desired result.

1 For the variant, it's the same except for a modified 0 clause:

$$2 \quad 0 \text{ iff } (\forall x \in W_w)[|A|_{x,v} = 1 \supset (\exists y \leq_w x)[|A|_{y,v} = 1 \wedge \\ 3 \quad (\forall z \leq_w y)(|A|_{z,v} = 1 \supset |B|_{z,v} = 0)]] \wedge (\exists x \in W_w)(|A|_{x,v} = 1).$$

4 Choose whichever you like: the construction that follows works with either choice.

5 To get the revision process started, we need a starting valuation j_0 , and we want it
6 to be transparent since this will guarantee that later j_v are as well. For simplicity I'll
7 take a trivial j_0 , which assigns value $\frac{1}{2}$ to each conditional at each world. It makes little
8 difference, because the effect of the starting values gets *almost* completely wiped out as the
9 construction proceeds. (It gets *completely* wiped out for sentences not containing 'True':
10 whatever the starting values, any such sentence gets the value that it gets in the 2-valued
11 worlds model for the corresponding version of Burgess semantics by stage n , where n is
12 the maximum depth to which ' \triangleright ' is embedded in the scope of other ' \triangleright 's in A ; and it keeps
13 that value at all subsequent stages. So from stage ω on, all 'True'-free sentences get "the
14 value they should", whatever the starting valuation.)

15 Finally, we need a policy on limit stages. Here the choice is important, and we choose
16 continuity with respect to 1 and 0. That is, if λ is a limit ordinal then for any world w and
17 any conditional $A \triangleright B$, j_λ assigns the conditional 1 at a world if and only if for some $\mu < \lambda$,
18 for every ordinal ν in the open interval (μ, λ) assigns the conditional value 1 at that world;
19 and similarly for 0. (So "irregularity arbitrarily close to λ " at a world as well as "constant
20 $\frac{1}{2}$ sufficiently close to λ " at that world lead to value $\frac{1}{2}$ at λ at that world.)

21 We can summarize these choices in a single definition. For the semantics based on the
22 modified Burgess, which I prefer, it's

$$23 \quad j_{w,\kappa}(A \triangleright B) \text{ is } \begin{cases} 1 & \text{if } (\exists \mu < \kappa)(\forall \nu \in [\mu, \kappa])(\forall x \in W_w)[|A|_{x,\nu} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y,\nu} = 1 \wedge (\forall z \leq_w y)(|A|_{z,\nu} = 1 \supset |B|_{z,\nu} = 1)]] \\ 0 & \text{if } (\exists \mu < \kappa)(\forall \nu \in [\mu, \kappa])(\forall x \in W_w)[|A|_{x,\nu} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y,\nu} = 1 \wedge (\forall z \leq_w y)(|A|_{z,\nu} = 1 \supset |B|_{z,\nu} = 0)]] \wedge \\ & (\exists x \in W_w)(|A|_{x,\nu} = 1)] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

24 ($[\mu, \kappa)$ is the half-open interval of ν such that $\mu \leq \nu < \kappa$.) For the semantics based on the
25 original Burgess, modify the 0 clause in the obvious way.

26 It's evident that on either variant, each j_κ is transparent if all preceding j_ν are transparent;
27 so by transfinite induction, all are transparent.

28 At each world, all 'True'-free sentences get the desired value (i.e. the one given in
29 the 2-valued model from which we started) by stage ω , and keep it at later stages. But
30 there is much greater irregularity for sentences containing 'True', due to the interaction
31 between 'True' and ' \triangleright '.²¹ In particular there is no fixed point. How then are we to select a
32 privileged j ?

33 The sequence of j_ν is a revision sequence in the sense of Gupta and Belnap 1993.
34 (The revision sequence depends on the model M_0 , as well as on the choice of Burgess
35 or modified-Burgess.) One well-known feature of revision sequences is that there are
36 evaluations j that appear arbitrarily late in the revision process; indeed, there are ordinals
 κ such that for any $\mu \geq \kappa$ and any ζ , there is a $\nu \geq \zeta$ such that $j_\nu = j_\mu$.²² Call any

²¹ The sentence itself needn't even contain ' \triangleright ' for the irregularity to occur, because the use of 'True'
typically makes other sentences relevant to the evaluation.

1 infinite such κ *final* (relative to model M_0),²³ and let FIN (or FIN_{M_0}) be the class of
2 final ordinals.

3 But not all final ordinals assign the same j (if they did, it would be a fixed point). Which
4 to pick? Obviously we want one that will yield as nice laws for \triangleright as possible. Gupta and
5 Belnap 1993 have a general theorem, their Reflection Theorem, that we can bring to bear.
6 Applied to this case, that theorem says:

7 PROPOSITION. [*Gupta-Belnap*] *There are limit ordinals Ω (“reflection ordinals for the*
8 *sequence j_κ ”) ²⁴ such that*

9 (i) Ω is *final*

10 (ii) For any L^+ -formulas A and B , and any world w and any $d \in \{0, \frac{1}{2}, 1\}$,

11 $(\exists \mu < \Omega)(\forall v \in [\mu, \Omega])(j_{w,v}(A \triangleright B) = d)$ if and only if $(\forall v \in FIN)$
12 $(j_{w,v}(A \triangleright B) = d)$.

13 Moreover, in the above semantics these reflection ordinals have an especially useful
14 property:

15 PROPOSITION. [*Fundamental Theorem for L (revision-theoretic version).*] For any
16 reflection ordinal Ω , any $w \in W$, and any L^+ -sentence A ,

17 (a) $|A|_{w,\Omega} = 1$ if and only if $(\forall v \in FIN)(|A|_{w,v} = 1)$

18 and (b) $|A|_{w,\Omega} = 0$ if and only if $(\forall v \in FIN)(|A|_{w,v} = 0)$.

19 Since there is only one possible value other than 0 and 1, these two clauses imply that
20 each reflection ordinal Ω is associated with the same j_Ω . This j_Ω is the valuation for \triangleright -
21 conditionals that I’ll be employing, e.g. in determining validity.

22 The Fundamental Theorem as stated here is similar to that given in Field 2008, but the
23 conditional there was different. The proof given there included a proof of [Gupta-Belnap],
24 since I was unaware of their theorem at the time. (Belated apologies to them for not being
25 able to give credit.) A proof of the Fundamental Theorem for the language of this paper,
26 now relying on [Gupta-Belnap] to save work, is given in Appendix A.

27 Note that when A is a conditional $B \triangleright C$, the 1-clause of the Fundamental Theorem
28 together with the evaluation rules for \triangleright yield that for any reflection ordinal Ω and $w \in W$,

29 **1-clause:** $|B \triangleright C|_{w,\Omega} = 1$ if and only if $(\forall v \in FIN)(\forall x \in W_w)(|B|_{x,v} = 1 \supset$
30 $(\exists y \leq_w x)(|B|_{y,v} = 1 \wedge (\forall z \leq_w y)(|B|_{z,v} = 1 \supset |C|_{z,v} = 1)))$.

31 Since $\Omega \in FIN$, this yields a necessary but not sufficient condition for $|B \triangleright C|_{w,\Omega} = 1$
32 that involves no ordinals other than Ω :

33 **1-clause Corollary:** If $|B \triangleright C|_{w,\Omega} = 1$ then $(\forall x \in W_w)(|B|_{x,\Omega} = 1 \supset (\exists y \leq_w x)(|B|_{y,\Omega}$
34 $= 1 \wedge (\forall z \leq_w y)(|B|_{z,\Omega} = 1 \supset |C|_{z,\Omega} = 1)))$.

35 That is, since we’ve chosen to use j_Ω for our final valuation: the 1-clause we’ve adopted
36 is strictly stronger than the 1-clause of the Burgess and modified-Burgess semantics.

²² Since the revision sequence here is *Markovian* in the sense that for any ordinals μ , κ and ν , if
 $j_\mu = j_\kappa$ then $j_{\mu+\nu} = j_{\kappa+\nu}$, we can simplify to: for any ζ , there is a $\nu \geq \zeta$ such that $j_\nu = j_\kappa$. If
this holds for κ in a Markovian sequence, it is bound to hold for any $\mu > \kappa$.

²³ It isn’t really necessary to demand infinitude explicitly, it’s entailed by the rest, as the reader can
easily prove using ‘True’-free sentences where ‘ \triangleright ’ is embedded to depth n for arbitrarily large n .

²⁴ Which ordinals are reflection ordinals will depend on the starting model M_0 .

1 But since all final ordinals are infinite, all ‘True’-free sentences receive the same value at
 2 all final ordinals; this means that for such B and C the ‘if...then’ in the corollary becomes
 3 an ‘if and only if’. In other words, we’re guaranteed that *the Burgess/modified-Burgess*
 4 *1-clause is retained for ‘True’-free sentences.*

5 Moreover, as long as we have Weak Centering at w , the 1-clause corollary yields the
 6 following for all B and C (not just the ‘True’-free ones):

7 **Modus Ponens for \triangleright :** If $|B \triangleright C|_{w,\Omega} = 1$ and $|B|_{w,\Omega} = 1$ then $|C|_{w,\Omega} = 1$.

8 (The label ‘Modus Ponens’ is really appropriate only if we have Weak Centering at all
 9 normal w .)

10 Something similar holds for the 0-clause, though the details depend on which version
 11 of the 0 clause one uses. In both cases, we get strictly stronger conditions than would be
 12 given by direct application of the Burgess or modified Burgess rules: e.g. for the semantics
 13 based on modified Burgess we get

14 If $|B \triangleright C|_{w,\Omega} = 0$ then $(\forall x \in W_w)[|B|_{x,\Omega} = 1 \supset (\exists y \leq_w x)[|B|_{y,\Omega} =$
 15 $1 \wedge (\forall z \leq_w y)(|B|_{z,\Omega} = 1 \supset |C|_{z,\Omega} = 0)] \wedge (\exists x \in W_w)(|B|_{x,\Omega} = 1).$

16 But again, when confined to ‘True’-free sentences the ‘if’ becomes an ‘if and only if’: *the*
 17 *Burgess or modified Burgess 0 clause is also retained for ‘True’-free sentences.*

18 (When w is weakly centered, the above yields

19 **0 Law for \triangleright :** If $|B \triangleright C|_{w,\Omega} = 0$ and $|B|_{w,\Omega} = 1$ then $|C|_{w,\Omega} = 0$ (and indeed, $|C|_{x,\Omega} = 0$
 20 whenever $x \sim_w w$),

21 which also strikes me as desirable but will play no role in what follows. Had we based the
 22 semantics on the original Burgess, we’d have needed that w be strongly centered to get this
 23 result.)

24 **4.2. Where are we?** For each starting arithmetically standard worlds model M_0 for the
 25 ‘True’-free fragment L_0 of L (with \triangleright evaluated either by the standard Burgess or variant
 26 Burgess rules), we have chosen a transparent j_Ω to evaluate all L^+ -conditionals at each
 27 world (including those containing embedded conditionals and/or ‘True’), and a T to evalu-
 28 ate truth-claims at each world. The worlds, and their division into normal and non-normal,
 29 are the same in the new model as in the old. (In particular, if the old contains no non-normal
 30 worlds, the new one won’t either.) The assignment of accessibility sets W_w and pre-orders
 31 \leq_w is also the same in the new model as in the old; so are the assignments of extensions
 32 to predicates at each world. And at each world, j_Ω assigns the same values to ‘True’-
 33 free conditionals (and hence ‘True’-free sentences more generally) as the original model
 34 on M_0 did. Finally, by the transparency of j and the features of the Kripke construction,
 35 the truth predicate is naive; and since the model is arithmetically standard, there can be
 36 no worry about using formulas with ‘True’ in the induction rule or validating generalities
 37 (e.g. composition rules) whose instances are valid.²⁵

²⁵ A feature of the model as described is that it is not value-functional: the value of $A \triangleright B$ at a world isn’t determined wholly by the values of A and B at it and other worlds. The reason is that all these values are values at a reflection Ω , and these depend on values at all non-reflection ordinals in FIN . But it isn’t hard to use what’s been done here to construct an enriched value space (along the lines of Field 2008, Section 17.1) in which we do have value-functionality: the value space for that will have infinitely many values, not linearly ordered. (The space is a set of functions from an initial segment of the ordinals to $\{0, \frac{1}{2}, 1\}$, where the length of the initial segment is the

1 The following are laws of this construction: by which I mean, schemas all of whose in-
 2 stances are valid (whatever structural conditions, such as Weak Centering at normal worlds,
 3 we decide on):

- 4 • $A \triangleright A$
- 5 • $[A \triangleright (B \wedge C)] \triangleleft \triangleright [(A \triangleright B) \wedge (A \triangleright C)]$
- 6 • $[(A \triangleright C) \wedge (B \triangleright C)] \triangleright [(A \vee B) \triangleright C]$
- 7 • $[A \triangleright (B \wedge C)] \triangleright [(A \wedge B) \triangleright C]$.

8 These are laws both when the evaluation rule for \triangleright is based on the original Burgess rule
 9 and when it is based on the modified rule: the 0 clause makes no difference. Indeed on both
 10 constructions they are all *strong laws*, by which I mean that their instances have value 1 at
 11 all worlds of every model, not just all normal worlds. That's important because it means
 12 that the result of prefixing any string of \Box s and \Diamond s to one of these is also a law. Related, it
 13 guarantees other "regular behavior", such as that we can strengthen antecedents in the laws.
 14 That is, even though we don't want and don't get that $Y \triangleright Z$ entails $X \wedge Y \triangleright Z$ for variably
 15 strict conditionals, still if $Y \triangleright Z$ is a strong law then so is $X \wedge Y \triangleright Z$ (even if X is true only at
 16 non-normal worlds). Similarly, if $X \triangleright Y$ and $Y \triangleright Z$ are strong laws then so is $X \triangleright Z$.²⁶ Proving
 17 that the bulleted schemas are strong laws is straightforward.²⁷ Note that since $\Box(A \triangleleft \triangleright A)$
 is valid, then by naivety so are $\Box(\text{True}(\langle A \rangle) \triangleleft \triangleright A)$ and $\Box(\neg \text{True}(\langle A \rangle) \triangleleft \triangleright \neg A)$, and hence

distance between successive reflection ordinals.) But for purposes of this paper there's no need
 for value-functionality.

²⁶ The proof that "antecedent strengthening" and transitivity are legitimate for strong laws uses the
 Fundamental Theorem as applied to \triangleright -sentences. Let W^* be the set of worlds that are n -accessible
 from worlds for some n . (On reasonable assumptions this will just be W , but the proof doesn't
 need this.) For antecedent strengthening, suppose that $Y \triangleright Z$ has value 1 at all worlds at reflection
 ordinals. Then it has value 1 at all worlds at all final ordinals, which means that at all final ordinals
 and all worlds in W^* , if Y has value 1 then so does Z ; and that includes all worlds where X has
 value 1. From this it's evident that $X \wedge Y \triangleright Z$ has value 1 at all worlds in W (even those not in W^* ,
 since only those in W^* are accessible to them) at all final ordinals, and in particular at reflection
 ordinals. The argument for transitivity is similar.

²⁷ The key observation for all of them is that for $|X \triangleright Y|_{w, \Omega}$ to be 1, it suffices that for all worlds
 w^* and all final ordinals v , if $|X|_{w^*, v} = 1$ then $|Y|_{w^*, v} = 1$. Given that, it's simply a matter
 of relativizing the proof that one would give for the Burgess-based semantics in the ground level
 language to a given v . For instance, for the right to left direction of the second listed law: Suppose
 that $|A \triangleright B|_{w^*, v} = |A \triangleright C|_{w^*, v} = 1$. Then for every x in W_{w^*} such that $|A|_{w^*, v} = 1$, there is a
 $y_1 \leq_{w^*} x$ such that

$$(a) |A|_{y_1, v} = 1 \wedge (\forall z \leq_{w^*} y_1)[|A|_{z, v} = 1 \supset |B|_{z, v} = 1],$$

and for every y_1 in W_{w^*} such that $|A|_{y_1} = 1$, there is a $y_2 \leq_{w^*} y_1$ such that

$$(b) |A|_{y_2, v} = 1 \wedge (\forall z \leq_{w^*} y_2)[|A|_{z, v} = 1 \supset |C|_{z, v} = 1].$$

Since \leq_{w^*} is a pre-order on W_{w^*} , (a) entails its analog (a*) where y_2 replaces y_1 ; and that with
 (b) yields

$$|A|_{y_2, v} = 1 \wedge (\forall z \leq_{w^*} y_2)[|A|_{z, v} = 1 \supset |B \wedge C|_{z, v} = 1],$$

which entails $|A \triangleright B \wedge C|_{w^*, v} = 1$.

(This proof and the proofs of the other laws just given doesn't depend on the use of a reflection
 ordinal for our evaluation: that should be no surprise, since the Fundamental Theorem shows that
 a single sentence can only have value 1 at reflection ordinals if it has value 1 at all final ordinals.
 Where the fact that validity requires preservation of value 1 only at reflection ordinals is important
 is for inferences from premises: e.g. Modus Ponens (assuming Weak Centering at normal worlds)
 and $A \wedge B \vDash A \triangleright B$ (assuming Strong).)

1 $\Box(\text{True}(\neg A) \triangleleft \triangleright \neg \text{True}(A))$. (And by the remarks at the end of Section 3, this means
 2 that we have a general composition principle for negation: for any sentence x , the negation
 3 of x is true if and only if x is not true.)

4 The fact that the above laws all hold in the construction with naive truth is interesting,
 5 because these are exactly the axiom schemas that Burgess uses in the quantifier-free case
 6 for the ‘True’-free fragment of the language. He gives a completeness proof there, for a
 7 system with these axioms, a necessitation rule, and the rule that for any string P of \Box s and
 8 \Diamond s, if $\models P\Box(A \equiv B)$ then $\models P[(A \triangleright C) \equiv (B \triangleright C)]$. The last rule is inappropriately
 9 weak in the 3-valued framework: we want a rule that has bite even when A and B aren’t
 10 bivalent. (An adequate replacement requires the additional conditional ‘ \rightarrow ’ soon to be
 11 introduced).²⁸ More generally, because the 3-valued background is weaker, the Burgess
 12 axiomatization doesn’t give a complete proof-procedure in the 3-valued context.²⁹ Still, I
 13 think that the fact that his axioms carry over unchanged is some indication that adding a
 14 naive truth predicate hasn’t seriously compromised the laws of ‘ \triangleright ’ (and once we add the
 15 ‘ \rightarrow ’ things will look even better).

16 In addition, we’ve seen that as long as we restrict the ground models to those with Weak
 17 Centering at normal worlds (as is required for Modus Ponens in the ground language),
 18 then Modus Ponens for \triangleright also holds in the expanded logic with ‘True’. (Some of the
 19 laws obtained in the 2-valued logic by adding restrictions on the \leq_w can only be carried
 20 over straightforwardly to the full logic with ‘True’ when stated using the aforementioned
 21 conditional ‘ \rightarrow ’ that generalizes the material conditional. We’ll turn to that conditional in
 22 Section 5.)

23 That’s the revision construction.

24 **4.3. The fixed point construction.** As I’ve mentioned, one can also give a fixed point
 25 construction that yields a rather similar outcome. Again consider valuation functions j that
 26 assign values in $\{0, \frac{1}{2}, 1\}$ to each pair of a world and a \triangleright -conditional; again we’re only
 27 interested in valuation functions that are transparent. The idea is to show that there is a set
 28 \mathbf{J} of transparent valuations, with a distinguished member j^* , where we have

29 PROPOSITION. [*Fundamental Theorem for L (fixed point version).*] For any $w \in W$,
 30 and any L^+ -sentence A ,

- 31 (a) $|A|_{w,j^*} = 1$ if and only if $(\forall h \in \mathbf{J})(|A|_{w,h} = 1)$
 32 (b) $|A|_{w,j^*} = 0$ if and only if $(\forall h \in \mathbf{J})(|A|_{w,h} = 0)$.

33 So j^* plays more or less the role that the j_Ω for reflection Ω play in the revision ap-
 34 proach, and \mathbf{J} plays more or less the role of the set of those j that occur arbitrarily late in the
 35 revision process (i.e. at ordinals in FIN). Here too, the various valuations j get a semantics
 36 whereby for any L^+ -sentences A and B and any world w , $j(w, A \triangleright B)$ is determined in a
 37 natural way from the values that valuations related to j give to B in worlds near w where A
 38 has value 1; and the semantics gives the values in the original model to L^+ -sentences not

²⁸ The best replacement is:

$$\text{For any string } P \text{ of } \Box \text{ and } \Diamond \text{, if } \models P\Box(A \leftrightarrow B) \text{ then } \models P[(A \triangleright C) \leftrightarrow (B \triangleright C)];$$

here ‘ \Rightarrow ’ is defined from ‘ \rightarrow ’ and strengthens it in a way to be discussed in Section 5, and ‘ \leftrightarrow ’
 and ‘ \Leftrightarrow ’ are defined from ‘ \rightarrow ’ and ‘ \Rightarrow ’ in the obvious ways. (The displayed law with mixed
 biconditionals entails the versions with two ‘ \leftrightarrow ’ and with two ‘ \Leftrightarrow ’.)

²⁹ Indeed, the fact that we’ve restricted to arithmetically standard models immediately rules out the
 possibility of a complete proof procedure.

1 containing ‘True’. To get the proper intersubstitutivity of logical equivalents, one needs to
 2 set up the semantics in a slightly non-obvious way. I sketch the construction in Appendix B;
 3 it is a generalization to variably strict conditionals of the one in Field 2014, and that paper
 4 will enable the reader to easily fill out the sketch in the Appendix.

5 (The basic idea of using a fixed point on a set of valuations was suggested in Yablo 2003;
 6 but Yablo’s procedure didn’t cut down the set of valuations quantified over in the semantics
 7 of each world nearly far enough—indeed, highly irregular valuations were included—
 8 and this led to extreme failure of intersubstitutivity of logical equivalents in embedded
 9 conditionals. Introducing chains in the manner of Appendix B seems to be the simplest
 10 acceptable way of accommodating Yablo’s basic insight.)³⁰

11 The remarks in Section 4.2 about the revision construction carry over to the fixed point
 12 construction virtually unchanged. In particular, the laws listed there are valid here too
 13 (again, with Modus Ponens as long as the original model has Weak Centering at normal
 14 worlds).

15 **§5. “Material-like” conditionals.** Many uses of ‘if ... then’ in English are captured
 16 reasonably well by a variably strict conditional like ‘ \triangleright ’, but some uses are more in line with
 17 a material conditional: in particular, the conditional used to restrict universal quantification
 18 is. “All A are B ” can’t be rendered as $\forall x(Ax \triangleright Bx)$: that’s too strong when ‘ \triangleright ’ is an ordinary
 19 indicative (or subjunctive) conditional. For instance, “Everyone who will be elected Presi-
 20 dent in 2016 is female” might be true but “For everyone x , if x is elected President in 2016
 21 then x is female” presumably isn’t: on the ordinary indicative reading, Jeb Bush and many
 22 others are counterexamples even if unelected. In a 2-valued context, we can represent “All
 23 A are B ” as $\forall x(Ax \supset Bx)$, where this is short for $\forall x(\neg Ax \vee Bx)$. But in a 3-valued context
 24 with restrictions on excluded middle, we can’t use a \supset defined in terms of \neg and \vee (at least
 25 if we want such schemas as “All A are A ” and “All A are either A or B ” to be logical laws);
 26 we need a new conditional ‘ \rightarrow ’ or ‘ \Rightarrow ’, that reduces to \supset for 2-valued sentences just as
 27 our ‘ \triangleright ’ reduces to the “classical” variably strict conditional.³¹ I find it plausible that this
 28 quantifier-restricting conditional is contraposable, but I needn’t insist on this: I will simply
 29 take ‘ \Rightarrow ’ to be a contraposable conditional and ‘ \rightarrow ’ to be a non-contraposable one, and we
 30 can leave open for now which of the two is to be used to define restricted quantification.
 31 There is no need for separate theories of ‘ \rightarrow ’ and ‘ \Rightarrow ’: we can take the basic conditional
 32 to be the non-contraposable ‘ \rightarrow ’, and define $A \Rightarrow B$ as $(A \rightarrow B) \wedge (\neg B \rightarrow \neg A)$, which
 33 ensures that ‘ \Rightarrow ’ is contraposable. The basic ‘ \rightarrow ’ and the derived ‘ \Rightarrow ’ have uses other than
 34 for restricting quantification: as observed in note 28, they are also needed for some of the

³⁰ Yablo’s paper also suggests the use of multiple Kripke fixed points for ‘True’ instead of the minimal ones; that idea can be employed with any of the constructions for ‘ \triangleright ’ in this section, both revision-theoretic and fixed point, and has what are arguably some advantages. For further discussion (in a revision-theoretic context with a different conditional), see Field 2008, Section 17.5. Again, it doesn’t matter to the issues of this paper whether one makes these modifications.

³¹ I should note that the notation used in this paper is almost the reverse of the notation in Field 2014. There, the material-like conditional used to restrict quantification (which was assumed contraposable) was symbolized as \blacktriangleright , and \triangleright was its non-contraposable generalization; whereas \rightarrow was used to symbolize a conditional with very much the flavor of the \triangleright used here, though it wasn’t based on a Stalnaker-Lewis-Pollock-Burgess multiple worlds semantics. Sorry for any confusion, but I think the new notation distinctly better.

An alternative to introducing a new conditional and defining universal restricted quantification in terms of it is to take a binary restricted quantifier $(\forall x \ni Ax)Bx$ as primitive. One can define ‘ \Rightarrow ’ (though not ‘ \rightarrow ’) from it, as well as the other way around.

1 laws of ‘ \triangleright ’ (and for these purposes, ‘ \rightarrow ’ as well as ‘ \Rightarrow ’ is required). But though I’ll take
2 ‘ \rightarrow ’ as basic, ‘ \Rightarrow ’ will be the primary focus, because at least in my own view, it is this
3 contraposable one that is ordinarily used to restrict universal quantification.

4 There are several options in the literature for such a conditional ‘ \rightarrow ’ (or a corresponding
5 contraposable ‘ \Rightarrow ’). Some of these are broadly like the revision-theoretic and fixed point
6 options for \triangleright given in Section 4; but a key difference is that the valuations at a single world
7 look only at other values at that same world.

8 For the moment let’s ignore the interaction between ‘ \rightarrow ’ and ‘ \triangleright ’, and focus on a lan-
9 guage L^* just like L except that it has ‘ \rightarrow ’ instead of ‘ \triangleright ’. A language with both ‘ \rightarrow ’
10 and ‘ \triangleright ’ is far more interesting, and will be treated in Section 6. That is what we’ll need
11 for a proper logic for restricted quantification in naive truth theory, a matter I’ll turn to in
12 Section 7. But for the moment, I look at L^* , which has ‘ \rightarrow ’ only.

13 L^* , like L , contains ‘True’; if it didn’t, and could be given a 2-valued semantics, we
14 could just define \rightarrow from \neg and \vee in the usual way. As before, the semantics for ‘True’
15 will be given by Kripkean constructions in which valuations v (analogous to the previous j)
16 for ‘ \rightarrow ’ at each world are held fixed; the real work then consists in the specification of an
17 appropriate valuation for ‘ \rightarrow ’ at each world.

18 A revision-theoretic construction of such a valuation for ‘ \Rightarrow ’ was given in Field 2008;
19 instead of what I called the “Official Conditional”, given in Ch. 16, I now prefer the “first
20 variation” given in Section 17.5, which modifies the 0 clause.³² And I want to adapt it to
21 the non-contraposable ‘ \rightarrow ’. Since L^* contains ‘ \square ’, we need to add a worlds parameter;
22 but the semantics for ‘ \rightarrow ’ is given world-by-world, unlike for ‘ \triangleright ’, and is thus considerably
23 simpler. It goes like this:

$$24 \quad |A \rightarrow B|_{w,\alpha} = \begin{cases} 1 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha))[|A|_{w,\gamma} = 1 \supset |B|_{w,\gamma} = 1] \\ 0 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha))[|A|_{w,\gamma} = 1 \wedge |B|_{w,\gamma} = 0] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

25 If we then define ‘ \Rightarrow ’ from ‘ \rightarrow ’ as above, we get something similar but with a strengthened
26 1-clause:

$$27 \quad |A \Rightarrow B|_{w,\alpha} = \begin{cases} 1 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha))[|A|_{w,\gamma} \leq |B|_{w,\gamma}] \\ 0 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha))[|A|_{w,\gamma} = 1 \wedge |B|_{w,\gamma} = 0] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

28 Like the earlier construction with ‘ \triangleright ’, this construction gives rise to a set of final ordinals
29 that include reflection ordinals Δ , and a Fundamental Theorem just like the previous:

30 PROPOSITION. [*Fundamental Theorem for L^* (revision-theoretic version).*] For any
31 reflection ordinal Δ , any $w \in W$, and any L^+ -sentence A ,

32 (a) $|A|_{w,\Delta} = 1$ if and only if $(\forall \gamma \in FIN)(|A|_{w,\gamma} = 1)$

33 (b) $|A|_{w,\Delta} = 0$ if and only if $(\forall \gamma \in FIN)(|A|_{w,\gamma} = 0)$.

34 It can be shown that if the ‘True’-free fragment L_0^* is 2-valued, \rightarrow and \Rightarrow are each
35 equivalent to the material conditional \supset on L_0^* . (If the ‘True’-free fragment L_0^* is 3-valued,
36 as it would be if we were to add \triangleright to the language and used the modified-Burgess-based
37 semantics, then \Rightarrow behaves on it like the Lukasiewicz 3-valued conditional, and \rightarrow like a
38 less familiar one.)

³² This switch yields a cleaner relation between $|A \Rightarrow B \wedge C|$ on the one hand and $|A \Rightarrow B|$ and
 $|A \Rightarrow C|$ on the other: see the end of this section. That in turn is important for restricted quantifier
law 4a* in Section 7.

1 As with \triangleright , only the valuations at reflection ordinals are relevant to validity: an inference
 2 is valid iff in all starting models and all worlds w in them and all reflection Δ , if the
 3 premises have value 1 at w and Δ then so does the conclusion.

4 Alternatively, we could adapt the fixed point semantics, to get a set \mathbf{R} of valuations
 5 u assigning values in $\{0, \frac{1}{2}, 1\}$ to each \rightarrow -conditional at each world, with privileged
 6 member v^* . Again, the semantics for non-privileged members of \mathbf{R} is given by a somewhat
 7 complicated chain construction analogous to that in Appendix B, but again it very much
 8 simplifies for v^* : we get

9 PROPOSITION. [*Fundamental Theorem for L^* (fixed point version).*] For any $w \in W$,
 10 and any L^{*+} -sentence A ,

11 (a) $|A|_{w,v^*} = 1$ if and only if $(\forall u \in \mathbf{R})(|A|_{w,u} = 1)$

12 (b) $|A|_{w,v^*} = 0$ if and only if $(\forall u \in \mathbf{R})(|A|_{w,u} = 0)$.

13 Only the special v^* is used in the definition of validity.³³

14 I note two consequences of the Fundamental Theorems for L^* :

15 **Modus Ponens for \rightarrow and \Rightarrow :** $A, A \rightarrow B \models B$ (and hence $A, A \Rightarrow B \models B$)

16 **Weak Equivalence of $\neg(A \rightarrow B)$ and $\neg(A \Rightarrow B)$ to $A \wedge \neg B$:** The inference from
 17 either $\neg(A \rightarrow B)$ or $\neg(A \Rightarrow B)$ to $A \wedge \neg B$ is valid, and so are the reverse inferences.

18 Why is the second one called “Weak” Equivalence? Two reasons: (a) While (in the revision
 19 version) $|\neg(A \rightarrow B)|_{w,\Delta}$ (or $|\neg(A \Rightarrow B)|_{w,\Delta}$) is 1 iff $|A \wedge \neg B|_{w,\Delta} = 1$, there is no
 20 analogous claim for 0. (b) Even for 1, the result holds only for reflection Δ , not for all
 21 final ordinals. (Similarly in the fixed point case: the equivalence holds only at v^* , not
 22 at all valuations in \mathbf{R} .) A consequence of (b) is that $\neg(A \rightarrow B)$ won’t in general be
 23 intersubstitutable with $A \wedge \neg B$ even in positive contexts, unless those contexts are outside
 24 the scope of \rightarrow ’s.

25 The proofs of Modus Ponens and Weak Equivalence are routine applications of the Fun-
 26 damental Theorem (for the appropriate construction) together with the evaluation clauses
 27 for \rightarrow . (Here there is no dependence on any Weak Centering assumption since the \rightarrow
 28 construction operates only within worlds.)

29 Later I will use the following (stated here for the revision-theoretic construction, but
 30 with analogs for fixed point): for all worlds w , and all ordinals α for (L-i) and all reflection
 31 ordinals Δ for (L-ii):

32 **(L-i):** If $|A \rightarrow B|_{w,\alpha} = 1$ then $|B \rightarrow C|_{w,\alpha} \leq |A \rightarrow C|_{w,\alpha}$;

33 **(L-ii):** $|A \rightarrow (B \wedge C)|_{w,\Delta} = \min\{|A \rightarrow B|_{w,\Delta}, |A \rightarrow C|_{w,\Delta}\}$.

34 The analogs for ‘ \Rightarrow ’ hold as well. Verification of (L-i) is almost trivial. (I’ll actually use it
 35 only in the case where α is a reflection ordinal, but it holds for all ordinals α .) Part of (L-ii)
 36 also generalizes to all ordinals:

37 **(L-iiia):** If $|A \rightarrow B|_{w,\alpha} = 1$ then $|A \rightarrow C|_{w,\alpha} \leq |A \rightarrow (B \wedge C)|_{w,\alpha}$

³³ The difference between the fixed point constructions for \rightarrow and for \triangleright comes in the way that chains
 of valuations generate valuations: instead of the association given in Appendix B, here when Z is
 a chain of \rightarrow -valuations we use the much simpler:

$$val[Z](w, A \rightarrow B) = \begin{cases} 1 & \text{if } (\exists S \in Z)(\forall u \in S)(|A|_{w,u} = 1 \supset |B|_{w,u} = 1) \\ 0 & \text{if } (\exists S \in Z)(\forall u \in S)(|A|_{w,u} = 1 \wedge |B|_{w,u} = 0) \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

This is basically what’s in Section 7 of Field 2014.

1 (and similarly for \Rightarrow), which is likewise easily proved. The remainder of (L-ii) is that
 2 when $|A \rightarrow (B \wedge C)|_{w,\Delta} = 0$, one of $|A \rightarrow B|_{w,\Delta}$ and $|A \rightarrow C|_{w,\Delta}$ must be 0. That's so
 3 because if $|A \rightarrow B|_{w,\Delta}$ and $|A \rightarrow C|_{w,\Delta}$ are both > 0 then (by the Fundamental Theorem
 4 and the evaluation rules) either there's a final α with $|A|_{w,\alpha} < 1$, or both a final α with
 5 $|B|_{w,\alpha} > 0$ and a final β with $|C|_{w,\beta} > 0$; and then by the Fundamental Theorem again,
 6 either $|A|_{w,\Delta} < 1$, or both $|B|_{w,\Delta} > 0$ and $|C|_{w,\Delta} > 0$. So $|A \rightarrow (B \wedge C)|_{w,\Delta+1} > 0$ and
 7 (by the Fundamental Theorem once again) $|A \rightarrow (B \wedge C)|_{w,\Delta} > 0$.³⁴

8 **§6. The two types of conditionals together.** So, we know several ways of getting
 9 naive truth in a language L with ' \triangleright ', and corresponding ways of getting naive truth in a
 10 language L^* with ' \rightarrow '. But what we really want is a language L^{**} with both (and with no
 11 restrictions on the embedding of either within the scope of the other).

12 There are three *prima facie* possible ways to proceed.

13 The *symmetric option* is to give a single construction (revision or fixed point, as one
 14 chooses) that evaluates both kinds of conditionals simultaneously: on the revision
 15 approach, this would involve, at each stage α , evaluating both $|A \triangleright B|_{w,\alpha}$ and $|A \rightarrow B|_{w,\alpha}$
 16 on the basis of the various $|A|_{x,\beta}$ and $|B|_{x,\beta}$ for $\beta < \alpha$ (restricting to the case where x is
 17 w in the case of \rightarrow).

18 The *\triangleright -first option* is to temporarily hold a valuation v for \rightarrow fixed, and use a construction
 19 for \triangleright on the basis of it. In the case of a revision construction, this would lead, for each
 20 choice of v , to a reflection ordinal Ω_v and thus a privileged valuation $j^v (= j_{\Omega_v}^v)$ for \triangleright ; in
 21 the case of a fixed point construction we similarly get a privileged valuation j^{*v} . Call this
 22 the "inner construction". We then would give an "outer" construction (again either revision-
 23 theoretic or fixed point; and it needn't be the same choice as for the inner) of a valuation for
 24 \rightarrow , one that looks only at the privileged valuations of \triangleright -conditionals constructed in inner
 25 constructions from other valuations. For instance, in the case where both the inner and outer
 26 constructions are revision-theoretic, we would construct $v_{\alpha+1}$ using valuations of sentences
 27 where \rightarrow -conditionals are evaluated by v_α and \triangleright -conditionals by the corresponding j^{v_α}
 28 (and use the same rule for limit ordinals as before), eventuating in a reflection ordinal Δ
 29 for the whole construction.

30 The *\rightarrow -first option* is just the reverse. In the case when both inner and outer constructions
 31 are revision-theoretic, we temporarily hold fixed a valuation j for \triangleright , and use a revision
 32 construction for \rightarrow on the basis of it; this leads, for each choice of j , to a reflection ordinal
 33 Δ_j and thus a privileged valuation $v^j (= v_{\Delta_j}^j)$ for \rightarrow . That is the "inner construction".
 34 We then would give an "outer" construction of a valuation j for \triangleright , where each $j_{\mu+1}$
 35 is determined from an evaluation of sentences that uses j_μ and the corresponding v^{j_μ} ,
 36 eventuating in a reflection ordinal Ω for the whole construction.

37 These three choices lead to significantly different results for the joint logic of \triangleright and \rightarrow .
 38 I think the \rightarrow -first option is most natural: very roughly, it involves settling the valuation
 39 of \rightarrow at each world before doing the \triangleright -construction which relates different worlds. But
 40 the ultimate rationale for the \rightarrow -first option is that it leads to by far the most plausible
 41 and useful laws of restricted quantification.³⁵ Some of the laws it leads to will be listed
 in Section 7. Few of them would hold on either the symmetric or \triangleright -first options: in the

³⁴ Had we used the valuation rules for the "Official Conditional" of Field 2008, we would only have gotten (L-ii), not (L-ii).

³⁵ Field 2014 used fixed point constructions rather than revision constructions for inner and outer, but the decision to take the restricted quantifier conditional as inner was the same there as here. (Recall from note 31 the confusing difference in notation: the restricted quantifier conditional

1 case of the revision construction, that's because on those options, the validity of a sentence
 2 of form $A \triangleright B$ (where A and B may contain \rightarrow) would require that B has value 1 when
 3 A does *at all final ordinals* in the \rightarrow -construction, not just at reflection ordinals of the
 4 \rightarrow -construction. For instance, it's only at reflection ordinals where A and $A \rightarrow \perp$ are
 5 prevented from simultaneously having value 1; because of this, the law $[(A \rightarrow B) \wedge A] \triangleright B$
 6 couldn't possibly hold on the symmetric or \triangleright -first options, where it does on the \rightarrow -first.
 7 (Similar remarks hold for the fixed point constructions.) For more remarks related to this,
 8 see note 42 below.

9 Let's recap (or make explicit) how the overall construction goes on the \rightarrow -first option.
 10 (I'll stick to the case where both the inner and outer constructions are revision-theoretic.)
 11 We start with a 2-valued worlds model M_0 for the 'True'-free fragment of L^{**} (whose
 12 number-theoretic part is an ω -model in each world, as before). Its ground fragment L^{**}_0
 13 is to be evaluated either by Burgess 2-valued or variant-Burgess 3-valued semantics. In the
 14 former case, ' \rightarrow ' is to be evaluated like ' \supset ' in the ground language. In the latter case, it is
 15 to be evaluated in the ground language by the rule that $|A \rightarrow B|$ is 1 whenever $|A| < 1$
 16 and is $|B|$ when $|A| = 1$. (This leads to \Rightarrow being evaluated in the ground language by the
 17 3-valued Lukasiewicz rules: $|A \Rightarrow B|$ is 1 iff $|A| \leq |B|$, 0 iff $|A|$ is 1 and $|B|$ is 0, $\frac{1}{2}$ iff $|A|$
 18 exceeds $|B|$ by $\frac{1}{2}$.) For convenience we expand the language L^{**} by adding names for all
 19 objects in the domain U of M_0 , getting L^{**+} .

20 Now let T be any function that assigns to every object of the ground model a value in
 21 $\{0, \frac{1}{2}, 1\}$, subject to the condition that if an object isn't the Gödel number of a sentence
 22 of L^{**} , T assigns it 0. Let j be any function that assigns to every L^{**+} -sentence of form
 23 $A \triangleright B$ a value in $\{0, \frac{1}{2}, 1\}$, and v be any function that assigns to every L^{**+} -sentence of
 24 form $A \rightarrow B$ a value in $\{0, \frac{1}{2}, 1\}$. We now evaluate every L^{**+} -sentence relative to T , j ,
 25 and v by essentially the Kleene rules early in Section 4; the only differences are that there
 26 is an additional parameter v in all the valuations, and we have an additional trivial clause
 27 for v analogous to that for j :

$$28 \quad |A \rightarrow B|_{w,j,v,T} = v(w, A \rightarrow B).$$

29 Then, keeping j and v fixed, we construct the minimal fixed point T_{min} (which now
 30 depends on v as well as on M_0 and j), and abbreviate $|A|_{w,j,v,T_{min}}$ as $|A|_{w,j,v}$.

31 Next we do the "inner construction": we hold the valuation j for \triangleright -sentences fixed, and
 32 do a revision construction for valuations v_α of \rightarrow -sentences. Adding a subscript j to make
 33 explicit the dependence on that \triangleright -valuation, the stages are given by:

$$34 \quad |A \rightarrow B|_{w,j,\alpha} = \begin{cases} 1 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha])(|A|_{w,j,\gamma} = 1 \supset |B|_{w,j,\gamma} = 1) \\ 0 & \text{if } (\exists \beta < \alpha)(\forall \gamma \in [\beta, \alpha])(|A|_{w,j,\gamma} = 1 \wedge |B|_{w,j,\gamma} = 0) \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

35 For each j , we are led to reflection ordinals Δ (which may depend on j as well as on
 36 M_0). And the dependence on a j clearly does nothing to block the Fundamental Theorem:
 37 we have

38 PROPOSITION. [*Fundamental Theorem for \rightarrow in L^{**} .*] For any j , any j -reflection
 39 ordinal Δ , any $w \in W$, and any L^+ -sentence A ,

40 (a) $|A|_{w,j,\Delta} = 1$ if and only if $(\forall \gamma \in FIN)(|A|_{w,j,\gamma} = 1)$

41 and (b) $|A|_{w,j,\Delta} = 0$ if and only if $(\forall \gamma \in FIN)(|A|_{w,j,\gamma} = 0)$.

42 Since there is only one possible value other than 0 and 1, these two clauses imply that
 each j -reflection ordinal Δ is associated with the same \rightarrow -valuation v^Δ ; we can call this

there was \triangleright , and the \rightarrow there was somewhat in the spirit of the \triangleright here.) The inner construction
 there was called the "fiber construction", and the outer construction the "base space construction".

1 valuation $v(j)$. Since the particular Δ doesn't matter as long as it is a j -reflection ordinal,
2 we can define $|A|_{w,j}$ to be $|A|_{w,j,\Delta}$ where Δ is any j -reflection ordinal.

3 In short, for each j -valuation for \triangleright -sentences, we've assigned a privileged valuation
4 $v(j)$ for \rightarrow -sentences. (And a minimal fixed point for truth, based on both.) That's the
5 inner construction.

6 We now use the privileged $v(j)$'s for each j in constructing a specific j for \triangleright -sentences
7 (the "outer construction"). So unlike in the inner construction, we don't need to add a
8 new parameter v for the valuation of the other conditional \rightarrow : the clauses for the j_μ that
9 evaluate \triangleright -sentences are EXACTLY as in Section 4.

10 This may seem to simplify matters, but it actually makes them somewhat more complicated:
11 for the v we use is no longer held constant, it varies with the j in the revision process.
12 Because of this, we need to revisit the Fundamental Theorem for \triangleright : in particular, the induction
13 on complexity in Stage (2) of the proof in Appendix A. For now we must consider in
14 the induction step not only sentences of form $\neg B$, $B \wedge C$, $\forall x B$ and $\Box A$, but also sentences
15 of form $A \rightarrow B$. And it's unobvious how to carry out the induction step in this case.

16 Indeed, it's more than unobvious: it can't be done, **the Fundamental Theorem for \triangleright**
17 **no longer holds without restriction once \rightarrow is added to the language.** Example: As a
18 preliminary, let K_\triangleright be constructed (by the usual Gödel-Tarski procedure) to be equivalent
19 to $True(\langle K_\triangleright \rangle) \triangleright \neg True(\langle K_\triangleright \rangle)$, and hence given naivety to $K_\triangleright \triangleright \neg K_\triangleright$. On the semantics
20 as given, at each world w for which $W_w \neq \emptyset$ (which includes all those w at which there is
21 at least Weak Centering) and for each stage κ for the outer construction,

22 **(\\$):** $|K_\triangleright|_{w,\kappa}$ is 1 if κ is odd, 0 if κ is an even successor, and $\frac{1}{2}$ if κ is a limit.

23 (That's so both for the semantics based on the original Burgess and the one based on
24 the variant.) Now let K^* be $K_\triangleright \rightarrow \neg K_\triangleright$. Since K_\triangleright is equivalent to a \triangleright -conditional, its
25 value is held fixed during any \rightarrow -construction, so at each w and each stage κ for the outer
26 construction and each stage $\alpha > 0$ for the inner, $|K^*|_{w,\kappa,\alpha}$ is 1 if $|K_\triangleright|_{w,\kappa} < 1$ and is 0
27 otherwise. So using (\$), when $W_w \neq \emptyset$, $|K^*|_{w,\kappa}$ (i.e. $|K^*|_{w,\kappa,\Delta}$) is 1 when κ is even
28 (including when it is a limit), and 0 otherwise. So for any world, at $\kappa = \Omega$, K^* has value 1,
29 but not at all final κ in the \triangleright -construction.³⁶

30 The failure of the Fundamental Theorem for \triangleright is not devastating, for we still get the
31 special case of it for \triangleright -conditionals, which is what is needed for many laws, such as
32 Modus Ponens (assuming Weak Centering for \triangleright). Indeed, we get more generally that the
33 Fundamental Theorem for \triangleright holds for every sentence A in which all occurrences of 'True'
34 and ' \rightarrow ' are inside the scope of an ' \triangleright '.

35 The special case of the Fundamental Theorem for \triangleright is enough to establish that all
36 reflection ordinals in the j_v construction give rise to the same values for every sentence:
37 for it immediately gives this for every \triangleright -conditional, and the generalization to all sentences
38 is immediate by induction.

39 **§7. Application to restricted quantification.** Here are some highly desirable laws of
40 restricted quantification: it is hard to imagine making serious use of restricted quantification
41 without them, or at least, something very close to them. Indeed, we should expect them

³⁶ It won't help to alter the starting point of the \rightarrow -construction, e.g. by making conditionals start
with value $\frac{1}{2}$ at some worlds but 1 at some and 0 at others. There are several reasons, but the
main one is that the evaluation of K_\triangleright would even out by stage ω , so that (\$) would still hold for
infinite κ .

1 to be strong laws in the sense explained in Section 4.1, which guarantees that prefixing
 2 any string of \Box s and \Diamond s to one of them is also to be a law, and that they remain valid
 3 however their antecedent is strengthened.³⁷ (The four with an asterisk are obtained using
 4 \triangleright -contraposition from their unasterisked counterparts;³⁸ but since \triangleright -contraposition isn't
 5 generally valid for variably strict conditionals they need to be stated separately. The ones
 6 marked 'b' result from the corresponding ones marked 'a' by a kind of quasi-contraposition
 7 which is also not generally valid for variably strict conditionals.) I've written these laws
 8 with \Rightarrow , reflecting my view that the conditional for restricted quantification is contrapos-
 9 able, but until we get to CQ, every law on the list would remain valid were \Rightarrow to be replaced
 10 with \rightarrow .

- 11 **1:** $[\forall x(Ax \Rightarrow Bx) \wedge Ay] \triangleright By$ "If all A are B, and y is A, then y is B"
 12 **2:** $\forall x Bx \triangleright \forall x(Ax \Rightarrow Bx)$ "If everything is B, then all A are B"
 13 **2*:** $\neg \forall x(Ax \Rightarrow Bx) \triangleright \neg \forall x Bx$ "If not all A are B, then not everything is B"
 14 **3a:** $\forall x(Ax \Rightarrow Bx) \wedge \forall x(Bx \Rightarrow Cx) \triangleright \forall x(Ax \Rightarrow Cx)$
 15 "If all A are B and all B are C then all A are C"
 16 **3b:** $\forall x(Ax \Rightarrow Bx) \wedge \neg \forall x(Ax \Rightarrow Cx) \triangleright \neg \forall x(Bx \Rightarrow Cx)$
 17 "If all A are B and not all A are C then not all B are C"
 18 **4a:** $\forall x(Ax \Rightarrow Bx) \wedge \forall x(Ax \Rightarrow Cx) \triangleright \forall x(Ax \Rightarrow Bx \wedge Cx)$
 19 "If all A are B and all A are C then all A are both B and C"
 20 **4b:** $\forall x(Ax \Rightarrow Bx) \wedge \neg \forall x(Ax \Rightarrow Bx \wedge Cx) \triangleright \neg \forall x(Ax \Rightarrow Cx)$
 21 "If all A are B and not all A are both B and C then not all A are C"
 22 **4a*:** $\neg \forall x(Ax \Rightarrow Bx \wedge Cx) \triangleright \neg \forall x(Ax \Rightarrow Bx) \vee \neg \forall x(Ax \Rightarrow Cx)$
 23 "If not all A are both B and C then either not all A are B or not all A are C"
 24 **5:** $\neg \forall x(Ax \Rightarrow Bx) \triangleright \exists x(Ax \wedge \neg Bx)$
 25 "If not all A are B, then something is both A and not B"
 26 **5*:** $\forall x(\neg Ax \vee Bx) \triangleright \forall x(Ax \Rightarrow Bx)$
 27 "If everything is either not-A or B, then all A are B" / "If nothing is both A
 28 and not-B, then all A are B"
 29 **6:** $\exists x(Ax \wedge \neg Bx) \triangleright \neg \forall x(Ax \Rightarrow Bx)$
 30 "If something is both A and not B, then not all A are B"
 31 **CQ:** $\forall x(Ax \Rightarrow Bx) \triangleright \forall x(\neg Bx \Rightarrow \neg Ax)$ "If all A are B then all not-B are not-A".
 32 **CQ*:** $\neg \forall x(Ax \Rightarrow Bx) \triangleright \neg \forall x(\neg Bx \Rightarrow \neg Ax)$
 33 "If not all A are B then not all not-B are not-A".

34 (There is a bit of redundancy in the list: 2* follows by obvious laws from 5, and 2 from 5*.)
 35 CQ and CQ* strike me as less *obviously* desirable than the earlier members of the list.
 36 However, CQ together with 1 and 2 respectively (and double negation laws in the case of 2)
 37 yield:

- 38 **1c:** $[\forall x(Cx \Rightarrow Dx) \wedge \neg Dy] \triangleright \neg Cy$ "If all C are D, and y is not D, then y is not C"
 39 **2c:** $\forall x \neg Cx \triangleright \forall x(Cx \Rightarrow Dx)$ "If nothing is C, then all C are D"

40 And these do seem to me obviously desirable; indeed, no less so than the laws 1 and 2 from
 41 which they were obtained. It's unobvious how to get a plausible theory that delivers 1c and
 42 2c without delivering CQ (and probably CQ*), which I take to provide support for the latter.

³⁷ Note that though the proof of the latter in note 26 relied on the Fundamental Theorem, it used it only for \triangleright -sentences, so it still holds when \rightarrow is in the language.

³⁸ With double negation laws (and re-lettering) in the case of CQ*.

1 Still, someone willing to give up 1c and 2c could use the results of this paper to validate
 2 the laws of restricted quantification preceding CQ with a restricted quantifier based on \rightarrow
 3 instead of \Rightarrow .

4 Despite the desirability of these laws, it is not entirely easy to give an account of
 5 conditionals in naive truth theory that validate them all (even without the modal prefixes).
 6 Indeed, prior to Field 2014, no published theory came close. But there are two precursors
 7 worth mentioning, Beall *et al* 2006 and Beall 2009. Both are in a paraconsistent framework,
 8 which means (given reasonable assumptions that they accept) that they can't accept a
 9 restricted-quantifier analog of law 2c, or even of its rule form. For if Cx means $x = x \wedge A$
 10 and Dx means $x = x \wedge B$ then even the rule version of 2c requires that $\neg A$ imply
 11 $A \mapsto B$ (where \mapsto is the paraconsistent restricted quantifier conditional); and then Modus
 12 Ponens yields Explosion. To deal with this, both precursors propose that the conditional
 13 that restricts quantification be non-contraposable,³⁹ i.e. they disallow even the rule form of
 14 CQ for \mapsto (and CQ*, given previous note). Myself, I'm not happy with the loss of 2c; but
 15 neither account does well with other laws either.

16 Beall *et al* 2006 made an important contribution in focusing on the need of a logic of
 17 restricted quantification and introducing the idea of using two separate conditionals for it.
 18 The paper didn't show, or even claim, that a naive truth theory could be added without
 19 triviality to the main logics it considers (those in their Section 6); but their discussion is
 20 explicitly motivated by the hope/belief that this is so. (One of the authors explicitly stated
 21 several years later that the question of non-triviality was open: see Beall 2009, p. 121.)
 22 Putting any worries about lack of non-triviality proof aside, the main issue is over the laws.
 23 The good news is that their framework validates their analogues of laws 2 and 4a (taking
 24 the analogues to have their noncontraposable \mapsto in place of my contraposable \Rightarrow , as well
 25 as their relevance conditional in place of my \triangleright); hence also 2* and 4a*, assuming the
 26 interpretation in note 39. The bad news is that it doesn't validate any of the others (though
 27 it does validate rule forms of some of them). Also, the validation of 2 and 2* depends very
 28 directly on their assumption of the validity of

29 **(?):** $A \triangleright B \models A \mapsto B$.

30 And (?) immediately rules out the analog of my law 1 (when naive truth, Modus Ponens
 31 for \mapsto , and reasonable quantifier laws are present). The reason is that given reasonable
 32 quantifier laws, law 1 requires $[(A \mapsto B) \wedge A] \triangleright B$; and then (?) delivers

33 **Pseudo Modus Ponens:** $[(A \mapsto B) \wedge A] \mapsto B$.

34 And it's well-known that this is inconsistent with genuine Modus Ponens for \mapsto (i.e.
 35 $(A \mapsto B) \wedge A \models B$) in a naive theory (assuming the standard structural rules for validity
 36 mentioned in note 3).⁴⁰ The centrality of (?) to the derivation of law 2 suggests that no
 37 simple modification of the account is likely to yield laws 1 and 2 together.

³⁹ Interestingly, they take their main conditional (a relevance conditional, their analog of my \triangleright)
 to obey a rule form of contraposition. (Beall 2009 very clearly does; Beall *et al* 2006 is slightly
 equivocal: see p. 595 middle.) I take this to mean that their main conditional isn't a good candidate
 for an account of the ordinary indicative conditional: see the Trump example in Section 2.

⁴⁰ In the logic I've been advocating (with Weak Centering assumed so as to get Modus Ponens), we
 do have

$$C \vee \neg C, C \triangleright B \models C \Rightarrow B;$$

but the need for the excluded middle premise is sufficient to prevent the paradox.

1 The second precursor is Beall 2009 (pp. 119-226). It also used two separate conditionals
 2 for the logic of restricted quantification. It suggests three different options for the logic,
 3 and unlike Beall *et al* 2006, shows each to be compatible with naive truth. All of them
 4 validate (?), so again it is immediate that law 1 can't be satisfied. The situation for laws is
 5 slightly worse than Beall *et al* 2006. Beall's first two options validate only 4a and 4a* from
 6 the list (though the weaker rule forms of some of the others are validated). His third option
 7 validates only 2 and 2*; indeed, its method of achieving 2 and 2* causes it to violate even
 8 the rule form of 4a.

9 Without going into detail, the main problem in both Beall *et al* 2006 and Beall 2009
 10 arises because (a) a certain kind of "abnormal" worlds are essential to these accounts
 11 (unlike the present account, where they are optional); (b) at these worlds, both conditionals
 12 are very badly behaved; and (c) the validity of $X \triangleright Y$ (using my notation for their relevance
 13 conditional) requires that it be true at all normal worlds, which in turn requires that at
 14 all worlds including abnormal ones, Y is true when X is. Collectively these make it very
 15 hard for reasonable \triangleright -statements with \mapsto -conditionals in their antecedents or consequents
 16 to come out valid. (An additional problem arises because of the way that these accounts
 17 handle negation, via a shift in worlds: this immediately rules out laws like 3b and 4b.)

18 Field 2014 used a very different framework, and did manage to validate the entire list;
 19 but the semantics it employed for \triangleright seemed *ad hoc*. (That paper did note some common-
 20 alities between its \triangleright and the ordinary indicative conditional, but also pointed out that the
 21 conditional reduced to the material conditional rather than the indicative conditional in
 22 'True'-free contexts.)⁴¹

23 But I now note that the entire list is also validated on the semantics of the present paper,
 24 with its independently motivated \triangleright . (We also get Modus Ponens for \triangleright , if we insist on Weak
 25 Centering at normal worlds in the base model, as I think we clearly should.)

26 The real work in establishing the laws on the list has nothing to do with the quantifiers,
 27 it's all in the relation among conditionals. The laws we need are the results of prefixing the
 28 following with strings of \Box s and \Diamond s:

- 29 **I:** $[(A \Rightarrow B) \wedge A] \triangleright B$ (for 1)
 30 **IIIa:** $(A \Rightarrow B) \wedge (B \Rightarrow C) \triangleright (A \Rightarrow C)$ (for 3a)
 31 **IIIb:** $(A \Rightarrow B) \wedge \neg(A \Rightarrow C) \triangleright \neg(B \Rightarrow C)$ (for 3b)
 32 **IVa:** $(A \Rightarrow B) \wedge (A \Rightarrow C) \triangleright (A \Rightarrow B \wedge C)$ (for 4a)
 33 **IVb:** $(A \Rightarrow B) \wedge \neg(A \Rightarrow B \wedge C) \triangleright \neg(A \Rightarrow C)$ (for 4b)
 34 **IVa*:** $\neg(A \Rightarrow B \wedge C) \triangleright \neg(A \Rightarrow B) \vee \neg(A \Rightarrow C)$ (for 4a*)
 35 **V:** $\neg(A \Rightarrow B) \triangleleft \triangleright (A \wedge \neg B)$ (for 5, 2* and 6)
 36 **V*:** $(\neg A \vee B) \triangleright (A \Rightarrow B)$ (for 5* and 2)
 37 **C:** $(A \Rightarrow B) \triangleleft \triangleright (\neg B \Rightarrow \neg A)$ (for CQ)
 38 **C*:** $\neg(A \Rightarrow B) \triangleleft \triangleright \neg(\neg B \Rightarrow \neg A)$ (for CQ*)

⁴¹ Despite its reducing to the material conditional, we can in retrospect see the conditional of Field 2014 as pretty much a degenerate case of the indicative conditional of the present paper. For the construction there started from a classical first order model, which can be seen as a degenerate Burgess model with only one world, weakly centered (which in the one-world case means simply "accessible from itself"). In that degenerate case, ' \triangleright ' obviously coincides with the material conditional in the ground model. (The conditional there still differed in a small respect from the degenerate case of the current construction: it utilized what I there called "dynamic Kripke constructions". I have dropped them here since they don't yield the results that we want once we clearly focus on extending the *ordinary indicative* conditional to a language with 'True'.)

1 C and C* are of course entirely trivial given the definition of \Rightarrow in terms of \rightarrow . For most
 2 of the others, the proof is almost immediate from what has already been said, especially at
 3 the end of Section 5. (The analogs of these latter laws for \rightarrow hold equally.) For note that
 4 to establish that a claim of form $P(X \triangleright Y)$ is valid, where P is any string of \Box s and \Diamond s,
 5 it suffices to show (in the revision-theoretic version; but the fixed point is analogous) that
 6 for all worlds w and all final κ of the \triangleright -construction, if $|X|_{w,\kappa} = 1$ then $|Y|_{w,\kappa} = 1$. In
 7 other words, that for all w and κ , and all κ -reflection ordinals Δ_κ of the \rightarrow -construction,
 8 if $|X|_{w,\kappa,\Delta_\kappa} = 1$ then $|Y|_{w,\kappa,\Delta_\kappa} = 1$. Given this, the proof of I is immediate from “Modus
 9 Ponens for \rightarrow and \Rightarrow ”, and V from “Weak equivalence of $\neg(A \rightarrow B)$ and $\neg(A \Rightarrow B)$ to
 10 $A \wedge \neg B$ ”. And IIIa and IIIb follow from the special case of (L-i) (end of Section 5) where
 11 α is Δ , and IVa, IVb and IVa* from (L-ii). As for V*, if $|\neg A \vee B|_{w,\kappa,\Delta_\kappa} = 1$ then either
 12 $|A|_{w,\kappa,\Delta_\kappa} = 0$ or $|B|_{w,\kappa,\Delta_\kappa} = 1$, and so by the Fundamental Theorem for \rightarrow , either for all
 13 κ -final α , $|A|_{w,\kappa,\alpha} = 0$ or else for all κ -final α , $|B|_{w,\kappa,\alpha} = 1$; in either case, for all κ -final
 14 α , $|A|_{w,\kappa,\alpha} \leq |B|_{w,\kappa,\alpha}$. From this it clearly follows that for all final α $|A \Rightarrow B|_{w,\kappa,\alpha} = 1$
 15 and hence in particular that $|A \Rightarrow B|_{w,\kappa,\Delta_\kappa} = 1$.

16 This only scratches the surface of the logic of the system,⁴² but it is not my purpose
 17 here to explore it at all systematically: my purpose is simply to show that it does easily
 18 lead to obvious laws of restricted quantification, which other approaches to conditionals in
 19 naive truth theory (other than the *ad hoc* one of Field 2014) haven’t come close to meeting.
 20 And I think that by basing the laws on an independently motivated account of indicative
 21 conditionals, the resulting theory is quite natural.

22 In particular, it’s worth emphasizing that the use of two distinct conditional operators
 23 (which is essential for the compatibility of the logic with naive truth, since if \rightarrow and \triangleright
 24 were identified then we’d have the disastrous (?)) is independently motivated: as I argue at
 25 the beginning of Section 5, we can see independently of the laws recently listed that the
 26 indicative conditional and the conditional for restricted quantification must be different.

27 **Thanks:** Harvey Lederman, Graham Priest and two anonymous referees made comments
 28 that have led to significant improvements.

29 **Appendix A: Proof of Fundamental Theorem for L (revision-theoretic version).**

30 Since $\Omega \in FIN$, the right to left of (a) and (b) in the Theorem are trivial. Contraposing
 31 the left to right and making the Kripke-stages σ explicit, what we need to establish is that
 32 for any reflection ordinal Ω and any L^+ -sentence A :

33 (a*) $(\forall w \in W)[\text{if } (\exists v \in FIN)(|A|_{w,v} < 1) \text{ then } \forall \sigma (|A|_{w,\Omega,\sigma} < 1)], \text{ and}$

34 (b*) $(\forall w \in W)[\text{if } (\exists v \in FIN)(|A|_{w,v} > 0) \text{ then } \forall \sigma (|A|_{w,\Omega,\sigma} > 0)].$

⁴² The reader will note that the schemas I’ve listed and proved are ones where there are no
 occurrences of \triangleright inside the scope of an \rightarrow (or an \Rightarrow). This is no accident: the \rightarrow -first construction
 makes it much easier for a schema in which \rightarrow is in the scope of \triangleright to be valid than for one where \triangleright
 is in the scope of \rightarrow to be valid. I think that schemas of the latter sort tend to be far less important
 than the former (recall the frequently-voiced claim that embeddings of indicative conditionals in
 the scope of other operators are hard to interpret); that is the main reason I went for an \rightarrow -first
 option. (I have however made no prohibitions on the well-formedness of embeddings of \triangleright inside
 the scope of \rightarrow ; and with any valid schema such as those listed, there are instances of the schema
 with arbitrarily complex chains of embeddings of \triangleright and \rightarrow .)

Despite what I’ve just said, there are important laws that depend on the embedding of ‘ \triangleright ’ in
 the scope of ‘ \rightarrow ’, but these are mostly meta-rules, whose legitimacy is not blocked by the \rightarrow -first
 option. A typical example is the meta-rule stated in note 28, whose proof is routine.

1 We establish (a*) and (b*) in three steps:

2 (1) In the special case when A is a conditional $B \triangleright C$, the value of σ makes no difference,
3 and by the fact that Ω is a limit ordinal and the evaluation rules for conditionals are
4 continuous with respect to 1 and 0 at limits, the claims are just:

5 (a*-s) $(\forall w \in W)[\text{if } (\exists v \in FIN)(j_{w,v}(B \triangleright C) < 1) \text{ then } (\forall \mu < \Omega)(\exists v \in [\mu, \Omega))$
6 $[j_{w,v}(B \triangleright C) < 1]]$;

7 (b*-s) $(\forall w \in W)[\text{if } (\exists v \in FIN)(j_{w,v}(B \triangleright C) > 0) \text{ then } (\forall \mu < \Omega)(\exists v \in [\mu, \Omega))$
8 $[j_{w,v}(B \triangleright C) > 0]]$.

9 But by (ii) of [Gupta-Belnap] these are trivial.

10 (2) Given (1), we can show (for any σ) that if (a*) and (b*) hold for the special case
11 where A is of form ‘ $True(c)$ ’ when c denotes the Gödel number of a sentence, then
12 they hold for all L^+ -sentences A . This is a routine induction on complexity, counting
13 \triangleright -sentences as of complexity 0 for the purposes of the induction: the claim is trivial for
14 all other atomic sentences since they keep the same value at every revision-stage v , and
15 the induction step for sentences $\neg B$, $B \wedge C$, $\forall x B$ and $\Box B$ is easy. For instance for \Box :

16 (a) Suppose that for some world w , $(\exists v \in FIN)(|\Box B|_{w,v} < 1)$. Then $(\exists v \in FIN)$
17 $(\exists y \in W_w)(|B|_{y,v} < 1)$; reversing the quantifier order and applying the induction hypoth-
18 esis, we get that for some $y \in W_w$, $|B|_{y,\Omega,\sigma} < 1$ (for any σ), and so $|\Box B|_{w,\Omega,\sigma} < 1$.

19 (b) Suppose that for some world w , $(\exists v \in FIN)(|\Box B|_{w,v} > 0)$. Then $(\exists v \in FIN)$
20 $(\forall y \in W_w)(|B|_{y,v} > 0)$; so certainly for all y in W_w , $(\exists v \in FIN)(|B|_{y,v} > 0)$, and by
21 the induction hypothesis for all y in W_w , $|B|_{y,\Omega,\sigma} > 0$ (for any σ); so $|\Box B|_{w,\Omega,\sigma} > 0$ for
22 any σ .

23 (3) It remains only to show that for all Kripke-stages σ and all c that denote Gödel
24 numbers of sentences, (a*) and (b*) hold for sentences of form ‘ $True(c)$ ’. But this is
25 trivial when $\sigma = 0$, since $|True(c)|_{w,\Omega,0}$ is always $\frac{1}{2}$. We now show that if it holds for
26 $\sigma = \tau$ then it holds for $\sigma = \tau + 1$. Suppose c denotes B . Then by the assumption about τ
27 and the result (2), we get

28 $(\forall w \in W)[\text{if } (\exists v \in FIN)(|B|_{w,v} < 1) \text{ then } |B|_{w,\Omega,\tau} < 1]$

29 and the analog with ‘ > 0 ’ instead of ‘ < 1 ’; which by the transparency of the j_v -valuations
30 and the Kripke construction gives

31 $(\forall w \in W)[\text{if } (\exists v \in FIN)(|True(c)|_{w,v} < 1) \text{ then } |B|_{w,\Omega,\tau} < 1]$

32 and its analog. But by the valuation rules, $|B|_{w,\Omega,\tau}$ is the same as $|True(c)|_{w,\Omega,\tau+1}$, so
33 the result is established. The case where σ is a limit ordinal is trivial: no sentence of form
34 ‘ $True(c)$ ’ first passes from $\frac{1}{2}$ to another value at a limit stage of the Kripke construction.

35 **Appendix B: The fixed point construction for L .** Again, a valuation function is a
36 function that assigns to each world and L^+ -conditional a value in $\{0, \frac{1}{2}, 1\}$.

37 Let a *chain* be a set P of nonempty sets of transparent valuation functions, meeting the
38 condition that if $S_1, S_2 \in P$ then either $S_1 \subseteq S_2$ or $S_2 \subseteq S_1$.

39 Given a chain P , define a valuation function $val[P]$ (“the valuation function generated
40 by P ”) as:

$$41 \quad val[P](w, A \triangleright B) \text{ is } \begin{cases} 1 & \text{if } (\exists S \in P)(\forall j \in S)(\forall x \in W_w)[|A|_{x,j} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y,j} = 1 \wedge (\forall z \leq_w y)(|A|_{z,j} = 1 \supset |B|_{z,j} = 1)]] \\ 0 & \text{if } (\exists S \in P)(\forall j \in S)[(\forall x \in W_w)[|A|_{x,j} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y,j} = 1 \wedge (\forall z \leq_w y)(|A|_{z,j} = 1 \supset |B|_{z,j} = 0)]] \wedge \\ & (\exists x \in W_w)(|A|_{x,j} = 1)] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

1 (This is for the semantics based on modified Burgess; for that based on original Burgess,
2 the modification of the 0 clause is obvious.) Clearly each $val[P]$ is transparent, given that
3 members of $\cup P$ are.

4 Let $P_1 \leq P_2$ mean that every member of P_1 has a subset that's a member of P_2 .
5 (Having small members makes a chain bigger.) Chains that are smaller in this ordering
6 generate weaker valuation functions: if $P_1 \leq P_2$ then for all w , $val_w[P_1] \leq_K val_w[P_2]$.
7 (That's simply because the 1 clause and 0 clause both have form " $(\exists S \in P)(\forall j \in S) \dots$ ".)

8 Define a sequence J_μ of sets of transparent valuation functions:

$$9 \quad J_\mu = \{val[P] : P \text{ is a chain and } (\forall \beta < \mu)(\exists S \in P)(S \subseteq J_\beta)\}.$$

10 For $\mu > 0$ an equivalent and perhaps more intuitive definition of J_μ is: $\{val[P] : P \text{ is a}$
11 $\text{non-empty chain and } (\forall \beta < \mu)(\forall S \in P)(S \subseteq J_\beta)\}$. This is more restrictive about the
12 chains, but it's easy to see that any valuation generated by one of the chains in the original
13 is generated also by one of the more restrictive ones.

14 If $\mu < \nu$, $J_\nu \subseteq J_\mu$, so obviously we eventually reach a fixed point \mathbf{J} . That would be
15 uninteresting if \mathbf{J} were empty, but it can be shown (following the model of Field 2014,
16 section 7) that $\mathbf{J} \neq \emptyset$. So letting \mathbf{P} be the set of \mathbf{J} -chains (chains whose members are all
17 subsets of \mathbf{J}) we'll have

$$18 \quad (\mathbf{FP}): \mathbf{J} = \{val[P] : P \in \mathbf{P}\}.$$

19 This sets up a one-many correspondence between the j in \mathbf{J} and the P in \mathbf{P} . (The members
20 of \mathbf{J} are the analogs in this construction of the valuation functions associated with ordinals
21 in FIN in the revision construction.)

22 The \leq -minimal chain is $\{\mathbf{J}\}$; let j^* be the valuation it generates, i.e. $val[\{\mathbf{J}\}]$. This is the
23 analog, in the fixed point construction, of the valuation function at reflection ordinals. We
24 have

$$25 \quad |A \triangleright B|_{w, j^*} = \begin{cases} 1 & \text{if } (\forall j \in \mathbf{J})(\forall x \in W_w)[|A|_{x, j} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y, j} = 1 \wedge (\forall z \leq_w y)(|A|_{z, j} = 1 \supset |B|_{z, j} = 1)]] \\ 0 & \text{if } (\forall j \in \mathbf{J})[(\forall x \in W_w)[|A|_{x, j} = 1 \supset (\exists y \leq_w x) \\ & [|A|_{y, j} = 1 \wedge (\forall z \leq_w y)(|A|_{z, j} = 1 \supset |B|_{z, j} = 0)]] \wedge \\ & (\exists x \in W_w)(|A|_{x, j} = 1)] \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

27 In this case the Fundamental Theorem, as stated in the text, concerns the special nature
28 of the valuation function at j^* . Its proof and the proof of the fixed point result are a simple
29 adaptation of that in Section 7 of Field 2014.

30 BIBLIOGRAPHY

- 31 Beall, J. C. (2009). *Spandrels of Truth*. Oxford: Oxford University Press.
32 Beall, J. C., Brady, R., Hazen, A., Priest, G., & Restall, G. (2006). Relevant Restricted
33 Quantification. *Journal of Philosophical Logic*, **35**, 587–598.
34 Burgess, J. (1981). Quick Completeness Proofs for some Logics of Conditionals. *Notre*
35 *Dame Journal of Formal Logic*, **22**, 76–84.
36 Field, H. (2008). *Saving Truth from Paradox*. Oxford: Oxford University Press.
37 Field, H. (2014). Naive Truth and Restricted quantification: Saving Truth a Whole Lot
38 Better. *Review of Symbolic Logic*, **7**, 147–91.
39 Gupta, A. and Belnap, N. (1993). *The Revision Theory of Truth*. Cambridge, MA:
40 MIT Press.

- 1 Kripke, S. (1975). Outline of a Theory of Truth. *Journal of Philosophy*, **72**, 690–716.
- 2 Lewis, D. (1974). *Counterfactuals*. Cambridge, MA: Harvard University Press.
- 3 McGee, V. (1985). A Counterexample to Modus Ponens. *Journal of Philosophy*, **82**,
- 4 462–71.
- 5 Pollock, J. (1976). The Possible Worlds Analysis of Counterfactuals. *Philosophical*
- 6 *Studies*, **29**, 469–76.
- 7 Restall, G. (2007). Curry’s Revenge: The Costs of Non-Classical Solutions to the
- 8 Paradoxes of Self-Reference. In Beall J. C., editor. *Revenge of the Liar*, Oxford: Oxford
- 9 University Press.
- 10 Stalnaker, R. (1968). A Theory of Conditionals. In Rescher, N., editor. *Studies in Logical*
- 11 *Theory*, Oxford: Blackwell Publishers.
- 12 Yablo, S. (2003). New Grounds for Naive Truth Theory. In Beall, J. C., editor, *Liars and*
- 13 *Heaps*, Oxford: Oxford University Press.

14 DEPARTMENT OF PHILOSOPHY
15 NEW YORK UNIVERSITY
16 5 WASHINGTON PLACE
17 NEW YORK, NY 10003
18 and
19 DEPARTMENT OF PHILOSOPHY
20 UNIVERSITY OF BIRMINGHAM
21 EDGBASTON, BIRMINGHAM
22 B15 2TT UK
23 *Email:* hartryfield@nyu.edu